


Home-based versus centre-based cardiac rehabilitation (Review)

Taylor RS, Dalal H, Jolly K, Moxham T, Zawada A



**THE COCHRANE
COLLABORATION®**

This is a reprint of a Cochrane review, prepared and maintained by The Cochrane Collaboration and published in *The Cochrane Library* 2010, Issue 6

<http://www.thecochranelibrary.com>



TABLE OF CONTENTS

HEADER	1
ABSTRACT	1
PLAIN LANGUAGE SUMMARY	2
BACKGROUND	2
OBJECTIVES	3
METHODS	3
RESULTS	5
Figure 1.	5
Figure 2.	7
Figure 3.	8
DISCUSSION	11
AUTHORS' CONCLUSIONS	12
ACKNOWLEDGEMENTS	12
REFERENCES	12
CHARACTERISTICS OF STUDIES	15
DATA AND ANALYSES	36
Analysis 1.1. Comparison 1 Exercise capacity, Outcome 1 Exercise capacity 3-12 month.	37
Analysis 1.2. Comparison 1 Exercise capacity, Outcome 2 Exercise capacity 12-24 month.	38
Analysis 2.1. Comparison 2 Blood Pressure [mm Hg], Outcome 1 Systolic BP 3-12 month.	39
Analysis 2.2. Comparison 2 Blood Pressure [mm Hg], Outcome 2 Diastolic BP 3-12 month.	40
Analysis 3.1. Comparison 3 Blood lipids [mmol/l], Outcome 1 Total cholesterol 3-12 month.	41
Analysis 3.2. Comparison 3 Blood lipids [mmol/l], Outcome 2 HDL cholesterol 3-12 month.	42
Analysis 3.3. Comparison 3 Blood lipids [mmol/l], Outcome 3 LDL-cholesterol 3-12 month.	43
Analysis 3.4. Comparison 3 Blood lipids [mmol/l], Outcome 4 Triglycerides 3-12 month.	44
Analysis 4.1. Comparison 4 Smoking, Outcome 1 Smoking 3-12 month.	45
Analysis 5.1. Comparison 5 Mortality, Outcome 1 Mortality.	46
Analysis 6.1. Comparison 6 Completers, Outcome 1 Completers.	47
ADDITIONAL TABLES	47
APPENDICES	53
WHAT'S NEW	62
HISTORY	62
CONTRIBUTIONS OF AUTHORS	62
DECLARATIONS OF INTEREST	62
SOURCES OF SUPPORT	63
DIFFERENCES BETWEEN PROTOCOL AND REVIEW	63
INDEX TERMS	63

[Intervention Review]

Home-based versus centre-based cardiac rehabilitation

Rod S Taylor¹, Hayes Dalal², Kate Jolly³, Tiffany Moxham¹, Anna Zawada⁴

¹PenTAG, Peninsula Medical School, University of Exeter, Exeter, UK. ²Primary Care, Peninsula Medical School, Exeter & Lower Lemon Street Surgery, Truro, UK. ³Department of Public Health and Epidemiology, University of Birmingham, Birmingham, UK. ⁴Agency for Health Technology Assessment, Warsaw, Poland

Contact address: Rod S Taylor, PenTAG, Peninsula Medical School, University of Exeter, Veysey Building, Salmon Pool Lane, Exeter, EX2 4SG, UK. rod.taylor@pms.ac.uk.

Editorial group: Cochrane Heart Group.

Publication status and date: Edited (no change to conclusions), published in Issue 6, 2010.

Review content assessed as up-to-date: 6 July 2008.

Citation: Taylor RS, Dalal H, Jolly K, Moxham T, Zawada A. Home-based versus centre-based cardiac rehabilitation. *Cochrane Database of Systematic Reviews* 2010, Issue 1. Art. No.: CD007130. DOI: 10.1002/14651858.CD007130.pub2.

Copyright © 2010 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

ABSTRACT

Background

The burden of cardiovascular disease world-wide is one of great concern to patients and health care agencies alike. Traditionally centre-based cardiac rehabilitation (CR) programmes are offered to individuals after cardiac events to aid recovery and prevent further cardiac illness. Home-based cardiac rehabilitation programmes have been introduced in an attempt to widen access and participation.

Objectives

To determine the effectiveness of home-based cardiac rehabilitation programmes compared with supervised centre-based cardiac rehabilitation on mortality and morbidity, health-related quality of life and modifiable cardiac risk factors in patients with coronary heart disease.

Search methods

We updated the search of a previous review by searching the Cochrane Central Register of Controlled Trials (CENTRAL) in *The Cochrane Library* (2007, Issue 4), MEDLINE, EMBASE and CINAHL from 2001 to January 2008. We checked reference lists and sought advice from experts. No language restrictions were applied.

Selection criteria

Randomised controlled trials (RCTs) that compared centre-based cardiac rehabilitation (e.g. hospital, gymnasium, sports centre) with home-based programmes, in adults with myocardial infarction, angina, heart failure or who had undergone revascularisation.

Data collection and analysis

Studies were selected independently by two reviewers, and data extracted by a single reviewer and checked by a second one. Authors were contacted where possible to obtain missing information.

Main results

Twelve studies (1,938 participants) met the inclusion criteria. The majority of studies recruited a lower risk patient following an acute myocardial infarction (MI) and revascularisation. There was no difference in outcomes of home- versus centre-based cardiac rehabilitation in mortality risk ratio (RR) was 1.31 (95% confidence interval (CI) 0.65 to 2.66), cardiac events, exercise capacity standardised mean difference (SMD) -0.11 (95% CI -0.35 to 0.13), as well as in modifiable risk factors (systolic blood pressure; diastolic

Home-based versus centre-based cardiac rehabilitation (Review)

Copyright © 2010 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

1

blood pressure; total cholesterol; HDL-cholesterol; LDL-cholesterol) or proportion of smokers at follow up or health-related quality of life. There was no consistent difference in the healthcare costs of the two forms of cardiac rehabilitation.

Authors' conclusions

Home- and centre-based cardiac rehabilitation appear to be equally effective in improving the clinical and health-related quality of life outcomes in acute MI and revascularisation patients. This finding, together with an absence of evidence of difference in healthcare costs between the two approaches, would support the extension of home-based cardiac rehabilitation programmes such as the Heart Manual to give patients a choice in line with their preferences, which may have an impact on uptake of cardiac rehabilitation in the individual case.

PLAIN LANGUAGE SUMMARY

Comparison of different modes of cardiac rehabilitation

Heart disease is one of the most common causes of premature death and ill health. Cardiac rehabilitation (CR) aims to restore people with heart disease to health through a combination of exercise with education and psychological support. Traditionally centre-based cardiac rehabilitation programmes (e.g. either based within a hospital, gymnasium or a sport centre setting) are offered to individuals after cardiac events, while home-based cardiac rehabilitation programmes have been introduced in an attempt to widen access and participation. The aim of this review has been to determine the effectiveness of home-based cardiac rehabilitation programmes compared with supervised centre-based cardiac rehabilitation.

The study population in the trials were mainly male with a mean age of 52-69 years. Study findings indicate that both home and hospital-based interventions are similar in their benefits on risk factors, health-related quality of life, death, clinical events and costs. There was some weak evidence to suggest that home-based interventions were associated with a higher level of adherence.

The limitations of the review are that the recruitment of the included trials was limited to stable coronary heart disease patients either following an acute-MI or revascularization, but no other cardiac populations, such as heart failure. There has been considerable diversity in the variety of centre-based and home-based cardiac rehabilitation interventions.

Related reviews, including four other Cochrane reviews, can be looked at for a fuller picture of a broader review and more conclusions about cardiac rehabilitation and the effectiveness of its specific contributant interventions and in CHD and heart failure populations.

BACKGROUND

Coronary heart disease (CHD) is a major cause of death and disability. Globally there were an estimated 7.22 million deaths from CHD in 2002 (WHO 2004). Although CHD mortality has decreased in many developed countries, with recent advances in treatment and prevention over half of these people are surviving (Allender 2008).

Cardiac rehabilitation (CR) is offered to individuals after cardiac events to aid recovery and prevent further cardiac illness. Cardiac rehabilitation has been shown to improve physical health, and decrease subsequent morbidity and mortality (Jolliffe 2001; Taylor 2004). Cardiac rehabilitation programmes typically achieve this through exercise, education, behaviour change, counselling and support and strategies that are aimed at targeting traditional risk factors for cardiovascular disease. Cardiac rehabilitation is an es-

sential part of the contemporary care of heart disease and is considered a priority in countries with high prevalence of CHD and heart failure (Balady 2007; Graham 2007; NICE 2007; Stone 2005).

Although the beneficial effects of cardiac rehabilitation have been shown, participation remains sub-optimal. One of the main reasons people give for not accepting the invitation to attend cardiac rehabilitation are difficulties in regularly attending sessions at their local hospital and reluctance to take part in group-based classes (Beswick 2004). Home-based cardiac rehabilitation programmes have been introduced in an attempt to widen access and participation. In the UK home-based cardiac rehabilitation with a self-help manual - the Heart Manual - supported by a nurse facilitator (Lewin 1992) is a popular method of rehabilitation. Figures from the National Audit for Cardiac Rehabilitation (NACR) indicate

that of the 199 sites in UK and Republic of Ireland that currently provide cardiac rehabilitation, 39 (19.6%) of these sites are currently providing the Heart Manual with some 13,000 copies to patients in UK each year (Heart Manual 2008). The Heart Manual has also been used in Italy, Canada, Australia and New Zealand (Heart Manual 2008).

The one systematic review (of randomised controlled trials) to date comparing home- and centre-based cardiac rehabilitation concluded that the outcomes of both settings were broadly similar (Jolly 2006). However, the authors commented that this conclusion was subject to uncertainty given both the poor quality of and small size of included trials. The total sample size across all trials was only 750 patients and excluded heart failure. We are aware of the recent completion of two large UK-based randomised controlled trials comparing home- and centre-based cardiac rehabilitation (Dalal 2007; Jolly 2007). This review aims to update the evidence base for home and centre-based cardiac rehabilitation using Cochrane review methodology.

OBJECTIVES

To determine the effectiveness of home-based cardiac rehabilitation programmes compared with supervised centre-based cardiac rehabilitation on mortality and morbidity, health-related quality of life and modifiable cardiac risk factors in patients with coronary heart disease.

METHODS

Criteria for considering studies for this review

Types of studies

Randomised controlled trials (individual or cluster level) either parallel group, cross-over or quasi-randomised design were eligible for inclusion. Systematic reviews and meta-analyses were identified as a source of additional randomised controlled trials.

Types of participants

The study population includes adults with MI, angina, or who had undergone revascularisation (coronary artery bypass grafting, percutaneous transluminal coronary angioplasty or coronary artery stent), or heart failure who have taken part, or been invited to take part, in cardiac rehabilitation.

Studies of participants with heart transplants and those implanted with either cardiac resynchronisation therapy (CRT) or implantable defibrillators (ICD) were excluded.

Types of interventions

Home-based cardiac rehabilitation is defined as a structured programme with clear objectives for the participants, including monitoring, follow up visits, letters or telephone calls from staff, or at least self-monitoring diaries. The comparison group is centre-based cardiac rehabilitation based in a variety of settings (e.g. hospital physiotherapy department, University gymnasium, community sports centre).

Types of outcome measures

Mortality (cardiac and overall), morbidity (reinfarction, revascularisation, cardiac associated hospitalisation), exercise capacity, modifiable coronary risk factors (smoking behaviour, blood lipid levels, blood pressure), health-related quality of life, adverse events (withdrawal from the exercise programme), health service utilisation or costs and cost effectiveness. During the review (and before any data analysis) it was decided that the outcome of adherence to intervention should be included, as it may have an influence on the values of the other outcomes.

Search methods for identification of studies

As this review forms part of a broader review, that includes updates of three other Cochrane systematic reviews addressing cardiac rehabilitation (Jolliffe 2001; Rees 2004a; Rees 2004b) and a new review (Davies 2008) of interventions for enhancing uptake and adherence to cardiac rehabilitation, a generic search strategy was undertaken.

Electronic searches

Randomised controlled trials were identified from a previously published systematic review (Jolly 2006). This list of studies was updated by searching the Cochrane Central Register of Controlled Trials (CENTRAL) in *The Cochrane Library* (2007, issue 4), MEDLINE (2001 to January 2008), EMBASE (2001 to January 2008), CINAHL (2001 to January 2008), and PsycINFO (2001 to January 2008), Health Technology Assessment (HTA) and the Database of Abstracts of Reviews of Effects (DARE) databases were searched via the NHS Centre for Reviews and Dissemination (CRD) website (2001 to January 2008). Conference Proceedings were searched on Web of Science: ISI Proceedings (2001 to January 2008).

A full search of CENTRAL was undertaken. All other searches were run from 2001 as this is the earliest date of searches for the previous Cochrane reviews on cardiac rehabilitation (Jolliffe 2001; Rees 2004a; Rees 2004b). This date overlaps the dates of the searches of the previous review on this topic (Jolly 2006).

Searches were limited to RCTs, systematic reviews, and meta-analysis and a filter applied to limit by humans. No language or other limitations were imposed. Consideration was given to variations

in terms used and spellings of terms in different countries so that studies were not missed by the search strategy because of such variations. Search strategies were designed with reference to those of the previous systematic review (Jolly 2006) and in accordance with Cochrane methods (Higgins 2008). See Appendix 1 for details of search strategies.

Searching other resources

Reference lists of all eligible trials and systematic reviews were searched for additional studies.

Data collection and analysis

Selection of studies

The titles and abstracts of studies identified were screened and clearly irrelevant studies discarded. The full-text reports of all potentially relevant randomised and quasi-randomised trials were obtained and assessed independently for eligibility, based on the defined inclusion criteria, by two reviewers (RST and Philippa Davies). Any disagreement was resolved by discussion and in those few occasions where uncertainty remained, the opinion of two further reviewers was taken (KJ, AZ).

Data extraction and management

A revised data extraction form was used to incorporate new additions on quality assessment in the Cochrane Handbook (Higgins 2008). Relevant data regarding inclusion criteria (study design, participants, interventions, and outcomes), risk of bias (randomisation, blinding, attrition and outcome reporting) and results were extracted. In cases where insufficient data were reported (e.g. method of randomisation, statistical methods) authors were contacted for further information. Data extraction was carried out by a single reviewer (AZ) and checked by a second reviewer (RST). Excluded studies and reasons for exclusion are detailed in [Characteristics of excluded studies](#). Where necessary authors of included studies were contacted for missing information.

Assessment of risk of bias in included studies

In accord with the recently updated Cochrane Handbook for Systematic Reviews (Higgins 2008) and RevMan update (RevMan 5), risk of bias was assessed in terms of the of quality of random sequence generation and allocation concealment, description of drop-outs and withdrawals (including analysis by intention-to-treat), blinding (participants, personnel and outcome assessment) and selective outcome reporting. In addition evidence was sought that the groups were balanced at baseline and that intention to treat analysis was undertaken. The risk of bias in eligible trials

was assessed by a single reviewer (AZ) and checked by a second reviewer (RST).

Data synthesis

Data were processed in accordance with the Cochrane Handbook for Systematic Reviews of Interventions (Higgins 2008). We sought outcome results at follow up and the focus of this review was the between-group difference in home- versus centre-based groups. For dichotomous variables relative risks and 95% confidence intervals (CI) were derived for each outcome. For continuous variables mean differences and 95% CI were calculated for each outcome.

Heterogeneity amongst included studies was explored qualitatively (by comparing the characteristics of included studies) and quantitatively (using the chi-squared test of heterogeneity and I^2 statistic). Where appropriate, the results from included studies were combined for each outcome to give an overall estimate of treatment effect. A fixed-effect meta-analysis was used except where statistical heterogeneity was identified, in which case a random-effects model was used.

Given the variety of exercise capacity measures reported, results for this outcome were expressed as a standardised mean difference (SMD). Otherwise continuous outcomes were pooled as weighted mean difference (WMD). For the purposes of meta-analysis, in the one study that reported continuous outcomes, findings by three age subcategories results were pooled in order to produce a single omnibus score for each group.

We had intended to use stratified meta-analysis and meta-regression to further explore heterogeneity and examine potential treatment effect modifiers. Given the small number of included trials such analyses were deemed inappropriately underpowered. However, subgroup analyses undertaken within trials were noted.

Sensitivity analysis was used to examine two areas of uncertainty in this review. Firstly, for exercise capacity, in addition to pooling all trials using SMD, the majority of trials that reported outcomes as metabolic equivalents (METs) were pooled using WMD. Secondly, because of the lack of detailed reporting there was some doubt that whether the study [Kassaaian 2000](#) was a true comparison between home- and hospital-based cardiac rehabilitation or rather a comparison of hospital-based cardiac rehabilitation versus usual care. All meta-analyses were undertaken with and without the inclusion of this trial.

[Marchionni 2003](#) reported outcomes for home- and centre-based group according to three patient age subgroups (i.e. 45-65, 66-75, >75 years). These data have been pooled to obtain a single overall outcome result for home- and centre-based groups. Gordon et al compared two home-based exercise groups: physician-supervised nurse-case-managed programme ([Gordon 2002 Supervised](#)) and community-based programme ([Gordon 2002 Community](#)) versus centre-based cardiac rehabilitation programme, while study of Miller et al reported results in subgroups by different time of inter-

vention: home versus centre-based brief - 11 weeks (Miller 1984 Brief) and home versus centre-based expanded - 26 weeks (Miller 1984 Expanded). For each of these two studies, outcome results are reported separately for both home versus centre comparison groups.

RESULTS


Description of studies

See: [Characteristics of included studies](#); [Characteristics of excluded studies](#).

Results of the search

Our update cross-cardiac rehabilitation review electronic searches yielded a total 11,561 titles. After reviewing titles and abstracts 13 full papers were retrieved for possible inclusion. Reviewing reference lists of all eligible publications identified four more publications for possible inclusion. After examining the full text, four papers were excluded and six studies (reported in 13 papers) were included. The systematic review Jolly 2006 identified six trials (reported in eight papers) all of which met the inclusion criteria of this review. Therefore in total 21 papers reporting on 12 studies were included in the review. The study selection process is summarised in the QUOROM flow diagram shown in Figure 1.

Figure 1.



Included studies

We included 12 trials (1,938 participants) comparing a home-based to a centre-based cardiac rehabilitation programme. Two

of the studies had three comparison arms and these have been analysed separately giving in total 14 comparisons. Six of these studies were included in the previous systematic review, [Jolly 2006](#) ([Arthur 2002](#); [Bell 1998](#); [Carlson 2000](#); [Marchionni 2003](#); [Miller 1984 Brief](#), [Miller 1984 Expanded](#); [Sparks 1993](#)). Six new trials were identified: two, whose publication dates predated the searches for [Jolly 2006](#), but were missed from the review ([Gordon 2002 Community](#), [Gordon 2002 Supervised](#); [Kassaiian 2000](#)), and four published since 2003 (the end search date of [Jolly 2006](#) ([Dalal 2007](#); [Daskapan 2005](#); [Jolly 2007](#); [Wu 2006](#)).

Three studies were UK-based ([Bell 1998](#); [Dalal 2007](#); [Jolly 2007](#)); four were based in US ([Carlson 2000](#); [Gordon et al \(Gordon 2002 Community](#); [Gordon 2002 Supervised](#)); [Miller et al \(Miller 1984 Expanded](#); [Miller 1984 Brief](#)); [Sparks 1993](#)) and one each in Canada ([Arthur 2002](#)), Turkey ([Daskapan 2005](#)), Italy ([Marchionni 2003](#)), Iran ([Kassaiian 2000](#)) and China ([Wu 2006](#)). Most studies reported outcomes up to 12-month post-randomisation. Only three studies reported longer-term follow up ([Jolly 2007](#) 24 months; [Marchionni 2003](#) 14 months; [Arthur 2002](#) 18 months). Eight studies compared comprehensive programmes (i.e. exercise plus the education and/or psychological management) while the remainder reported only an exercise intervention ([Daskapan 2005](#); [Kassaiian 2000](#); [Miller 1984 Brief](#); [Miller 1984 Expanded](#); [Wu 2006](#)). The cardiac rehabilitation programmes differed considerably in duration (range: 1.5 to 6 months), frequency (1 to 5 sessions per week) and session length (20-60 minutes per session). Most programmes used individually tailored exercise prescription which makes it difficult to precisely quantify the amount of exercise undertaken. Centre-based programmes typically provided supervised cycle and treadmill exercise while virtually all

home programmes were based on walking, with some level of intermittent nurse or exercise specialist telephone support. The majority of studies recruited a lower risk patient following an acute MI and revascularisation, excluding those with significant arrhythmias, ischaemia, or heart failure. Two studies included individuals with New York Heart Association (NYHA) class 2 or 3 heart failure ([Daskapan 2005](#); [Kassaiian 2000](#)).

Details of included studies are listed in [Characteristics of included studies](#).

Excluded studies

Four papers were excluded: one was a comparison of two forms of home-based cardiac rehabilitation ([Senuzun 2006](#)), two were comparisons of home based cardiac rehabilitation versus usual care ([Sinclair 2005](#); [Tygesen 2001](#)) and one was a non-RCT ([Ades 2000](#)). Details of excluded studies are listed in [Characteristics of excluded studies](#).

Risk of bias in included studies

A number of studies failed to give sufficient detail to assess their potential risk of bias ([Figure 2](#)). Details of generation and concealment of random allocation sequence were particularly reported. In one case there was objective evidence of imbalance in baseline characteristics ([Arthur 2002](#)). Blinding of patients and carers in studies on CR is impossible; in such situations, blinding outcome assessment to knowledge of allocation may be of great importance. However only six of the studies stated that they took measures to blind outcome assessment, this may have weakened their conclusions.

Figure 2. Methodological quality summary: review authors' judgements about each methodological quality item for each included study.


	Adequate sequence generation?	Allocation concealment?	Blinding?	Incomplete outcome data addressed?	Free of selective reporting?	Groups balanced at baseline?	Intention to treat analysis?	Groups received same intervention?
Arthur 2002	?	+	+	+	+	-	+	+
Bell 1998	?	+	+	?	+	+	-	-
Carlson 2000	?	?	?	-	+	+	+	-
Dalal 2007	+	+	+	+	+	+	+	+
Daskapan 2005	?	?	?	+	+	+	-	+
Gordon 2002 Community	?	?	?	+	+	+	?	+
Gordon 2002 Supervised	?	?	?	+	+	+	?	+
Jolly 2007	+	+	+	+	+	+	+	-
Kassaian 2000	?	?	?	?	?	+	?	?
Marchionni 2003	?	?	+	+	+	+	+	+
Miller 1984 Brief	?	?	?	+	+	?	+	+
Miller 1984 Expanded	?	?	?	+	+	?	+	+
Sparks 1993	?	?	?	+	+	+	+	+
Wu 2006	?	?	+	?	+	+	?	+

Where reported, losses to follow up varied considerably across studies and was often asymmetric across home- and centre-based cardiac rehabilitation groups. Although often not stated, all studies appeared to undertake an intention to treat analysis in that groups were analysed according to initial random allocation. The impact of losses to follow up or drop out was only examined in a few trials. As discussed above, the rehabilitation intervention was usually tailored to the individual patient and therefore it is difficult to quantify the precise level of intervention. However, based on the general description of the intervention reported by authors, there appeared to be substantive differences in the nature of the rehabilitation input between home- and centre-based arm. For example,

the studies [Bell 1998](#), [Dalal 2007](#) and [Jolly 2007](#) included hospital cardiac rehabilitation programmes which were fixed in terms frequency and content over the period of the study. In contrast the home-based intervention in these studies consisted of the Heart Manual where the patients could self-regulate the frequency and nature of rehabilitation sessions they undertook.

Methodological quality graph ([Figure 3](#)) presents review authors' judgements about each methodological quality item presented as percentages across all included studies. Given the small number of included trials we were unable to assess publication bias using Funnel plot approach ([Higgins 2008](#)).

Figure 3. Methodological quality graph: review authors' judgements about each methodological quality item presented as percentages across all included studies.



Effects of interventions

Exercise capacity

All 12 included studies reported exercise capacity in the short-term (3 to 12 months follow up), while three ([Arthur 2002](#); [Jolly 2007](#); [Marchionni 2003](#)) presented longer-term data (12 to 24 months follow up). All reported exercise capacity at follow up, except one ([Gordon 2002 Supervised](#); [Gordon 2002 Community](#)) which instead reported change in exercise capacity at follow up

compared to baseline.

Nine studies reported exercise capacity as maximal oxygen uptake (VO_2max) either as metabolic equivalents (METs) or millilitres per kilogram of body mass per minute (ml/kg/min) or millilitres (ml) ([Sparks 1993](#)). [Jolly \(Jolly 2007\)](#) reported incremental shuttle walking distance (in metres, m), [Marchionni 2003](#) reported total cycle work capacity (in kilograms of body mass multiplied by meters ($kg \cdot m$)) and [Gordon \(Gordon 2002 Community](#); [Gordon 2002 Supervised](#)) reported change data only as METs.

In the pooled analysis across all studies reporting short-term data (1,938 patients) there was no evidence of a statistically significant difference in short-term exercise capacity between home-based and centre-based cardiac rehabilitation (random effects SMD = -0.11, 95% CI -0.35 to 0.13, [Analysis 1.1](#)). There was evidence of substantial heterogeneity ($I^2 = 79\%$; $\text{Chi}^2 = 60.91$; $P < 0.00001$). The same findings were seen when pooling was limited to the eight trials reporting VO_2max (random effects WMD = -0.30, 95% CI -1.22 to 0.63; $I^2 = 88\%$; $\text{Chi}^2 = 68.6$; $P < 0.0001$). Excluding [Kassaian 2000](#) from the later analysis reduced the degree of heterogeneity but produced the same finding of no evidence of significant difference between group (random effects WMD = 0.16, 95% CI -0.44 to 0.77, $P = 0.59$; $I^2 = 69\%$; $\text{Chi}^2 = 22.2$; $P = 0.002$).

In a pooled analysis of three studies ([Arthur 2002](#); [Jolly 2007](#); [Marchionni 2003](#)) reporting longer-term data (12 to 24 months; 1,074 patients) there was some evidence, albeit not statistically significant, of superior exercise capacity in the home- compared to the centre-based group (fixed effect SMD 0.11, 95% CI -0.01 to 0.23; $I^2 = 0\%$; $\text{Chi}^2 = 0.97$; $P = 0.62$; [Analysis 1.2](#)).

In all studies except two ([Dalal 2007](#); [Jolly 2007](#)) exercise capacity was assessed at baseline. In the remaining studies there was consistent evidence of increase in exercise capacity at follow up compared to baseline for both home- and centre-based groups.

Modifiable risk factors

Blood pressure

Seven of the included trials reported both systolic blood pressure (SBP) and diastolic blood pressure (DBP) ([Carlson 2000](#); [Dalal 2007](#); [Daskapan 2005](#); [Gordon 2002 Community](#); [Gordon 2002 Supervised](#); [Jolly 2007](#); [Kassaian 2000](#)), or SBP alone ([Bell 1998](#)). All studies reported outcome at follow-up, with the exception of one ([Gordon 2002 Supervised](#); [Gordon 2002 Community](#)) that instead reported change from baseline. For Dalal ([Dalal 2007](#)) non-published follow-up values were obtained on request from the authors. Blood pressure (BP) has been reported as millimetres of Hg (mmHg).

At 3 to 12 months follow up, although no between-group difference was found in pooled systolic blood pressure (random effects WMD = -0.58 mmHg, 95% CI -3.29 to 4.44; $I^2 = 70\%$; $\text{Chi}^2 = 23.01$; $P = 0.002$; [Analysis 2.1](#); 1,053 patients), there was a slightly higher pooled diastolic BP at follow up for home-based compared to centre-based cardiac rehabilitation (fixed effect WMD = 1.85 mmHg; 95%CI 0.74 to 2.96; $I^2 = 25\%$; $\text{Chi}^2 = 7.97$; $P = 0.24$; 927 patients; [Analysis 2.2](#)). However, this difference is not clinically relevant. On excluding [Kassaian 2000](#) this difference in DBP was no longer statistically significant (fixed effect WMD 1.00 mmHg; 95%CI -0.32 to 2.31; $I^2 = 0\%$; $\text{Chi}^2 = 2.28$; $P = 0.81$) while the finding of no difference in SBP remained (fixed effect WMD = -

1.06 mmHg, 95%CI -3.40 to 1.31; $I^2 = 32\%$; $\text{Chi}^2 = 8.76$; $P = 0.19$). At 24-months follow up, [Jolly 2007](#) reported no significant difference in SBP (mean 0.85 mmHg; 95%CI -2.48 to 4.18), or DBP (mean 0.76 mmHg, 95% CI -1.12 to 2.64) between home- and centre-based CR groups. There was no consistent trend across studies in change in blood pressure at follow up compared to baseline.

Blood lipids

Seven of the included trials reported data on blood lipids ([Bell 1998](#); [Carlson 2000](#); [Dalal 2007](#); [Gordon 2002 Community](#); [Gordon 2002 Supervised](#); [Jolly 2007](#); [Kassaian 2000](#)), all reporting total cholesterol values, four ([Carlson 2000](#); [Gordon 2002 Community](#); [Gordon 2002 Supervised](#); [Jolly 2007](#); [Kassaian 2000](#)) reporting HDL-cholesterol, and three ([Carlson 2000](#); [Gordon 2002 Community](#); [Gordon 2002 Supervised](#); [Kassaian 2000](#)) reporting LDL-cholesterol and triglycerides values. All studies but [Gordon 2002](#) reported follow up data, while [Gordon 2002 Community](#); [Gordon 2002 Supervised](#) only reported data for change. Study results were expressed as millimols per litre (mmol/l) ([Bell 1998](#); [Dalal 2007](#); [Jolly 2007](#)) or milligrams per decilitre (mg/dl) ([Carlson 2000](#); [Gordon 2002 Community](#); [Gordon 2002 Supervised](#); [Kassaian 2000](#)); in the latter case there have been converted into mmol/l before being pooled in the metaanalysis.

In all studies with the exception of [Kassaian 2000](#), compared to baseline there was evidence of a decrease in total cholesterol, LDL-cholesterol, and triglyceride levels, and increase of HDL-cholesterol levels at follow up in both home- and centre-based groups.

Total cholesterol

In the pooling analysis at 3 to 12 months follow up there was no evidence of a significant difference in the total cholesterol for the home and centre groups (random effects WMD = 0.13 mmol/l, 95% CI -0.05 to 0.31; $I^2 = 55\%$; $\text{Chi}^2 = 13.33$; $P = 0.04$; 1,019 patients; [Analysis 3.1](#)).

[Jolly 2007](#) reported no significant difference between home- and centre-based cardiac rehabilitation groups in total cholesterol (mean = 0.11 mmol/l, 95% CI: -0.06 to 0.28) at 24-months follow up.

HDL-cholesterol

In the pooling analysis at 3-12 months follow up there was evidence of a lower HDL-cholesterol for the home than for centre (fixed effect WMD = -0.06 mmol/l, 95% CI -0.11 to -0.02; $I^2 = 38\%$; $\text{Chi}^2 = 6.5$; $P = 0.16$; 793 patients; [Analysis 3.2](#)). When [Kassaian 2000](#) was excluded, there was no longer a significant difference between groups in HDL-cholesterol (fixed effect WMD

= -0.02 mmol/l, 95% CI -0.08 to 0.04; $I^2 = 0\%$; $\text{Chi}^2 = 0.5$; $P = 0.91$).

Jolly 2007 reported no significant difference between home- and centre-based cardiac rehabilitation groups in HDL-cholesterol (mean = 0.03 mmol/l, 95%CI -0.10 to 0.04) at 24-months follow up.

LDL-cholesterol

In the pooling analysis at 3-12 months follow up there was no evidence of difference in LDL-cholesterol (fixed effect WMD = 0.15 mmol/l, 95% CI -0.01 to 0.31; $I^2 = 32\%$; $\text{Chi}^2 = 4.4$; $P = 0.22$; 324 patients; Analysis 3.3). When Kassaian 2000 was excluded between group results remained non significant.

Triglycerides

In the pooling analysis at 3-12 months follow up there was no evidence of difference in triglycerides (random effects WMD = 0.15 mmol/l, 95% CI -0.11 to 0.41; $I^2 = 60\%$; $\text{Chi}^2 = 7.58$; $P = 0.06$; 328 patients; Analysis 3.4). When Kassaian 2000 was excluded between group results remained non significant.

Smoking behaviour

Five of the 12 studies included reported patient self-reported smoking behaviour at 3-12 months follow up (Bell 1998; Dalal 2007; Gordon 2002 Community; Gordon 2002 Supervised; Jolly 2007). There was no evidence of difference in the proportion of smokers at follow up with centre- and home-based cardiac rehabilitation (fixed effect RR = 1.00, 95% CI 0.71 to 1.41; $I^2 = 11\%$; $\text{Chi}^2 = 4.48$; $P = 0.34$; 922 patients; Analysis 4.1). Jolly 2007 reported no difference in smoking between home- and centre-based arms at 24-months (RR = 1.16, 95% CI 0.58 to 33.3).

There was evidence of a consistent reduction in self-reported smoking behaviour following both home- and centre-based cardiac rehabilitation. This finding was confirmed in the one study that used cotinine-validated assessment of smoking (Jolly 2007).

Health-related quality of life

Five out of the 12 included trials reported validated health-related quality of life (HRQoL) measures (Table 1). These included four generic HRQoL instruments EQ-5D (EuroQoL 1990), Nottingham Health Profile (Hunt 1980), Short-Form 36 (SF-36; McHorney 1993), Sickness Impact Profile (Bergner 1976) and one disease-specific instrument (MacNew; Höfer 2004). Given the wide variation in HRQoL outcomes used, pooling across studies was deemed inappropriate. The HRQoL results at follow up and between-group difference for each individual trials were not reported in the original publications and between-group P-values were calculated by the authors of this report using methods outlined in the Cochrane Handbook (Higgins 2008).

Overall there was no evidence of a statistically significant difference in overall HRQoL or domain score at follow up between home and centre-based groups. The two exceptions were a higher Nottingham Health Profile sleep domain score in the hospital compared to the home group in Bell 1998 and a higher SF-36 physical component score in the home compared to centre-based cardiac rehabilitation groups at 6 months in Arthur 2002.

Individual studies reported consistent improvements in HRQoL at follow up with both home and centre-based cardiac rehabilitation compared to baseline. The notable exception was the use of the EQ-5D which failed to identify significant improvements with home- or centre-based cardiac rehabilitation (Dalal 2007; Jolly 2007).

Clinical events

Mortality

Five trials reported all-cause mortality (specific cardiac death have not been separately reported in the included studies) in the period up to 1-year follow up (Bell 1998; Dalal 2007; Daskapan 2005; Jolly 2007; Miller 1984 Brief; Miller 1984 Expanded). The study of Miller (Miller 1984 Brief; Miller 1984 Expanded) reported no deaths in either the home- or centre-based CR groups over the period of the study (and so it has not been included in pooled analysis). In a pooled analysis of the remaining studies (909 patients) there was no evidence of a significant difference in mortality at 3-12 months follow up between home and centre (fixed effect RR = 1.31, 95% CI 0.65 to 2.66; $I^2 = 0\%$; $\text{Chi}^2 = 1.0$, $P = 0.8$; Analysis 5.1). Excluding Daskapan 2005 (because of patients with NYHA class 2/3 heart failure) the pooled result of 9-12 months studies shows no significant difference as well (RR = 1.25, 95% CI 0.61 to 2.59; $I^2 = 0\%$; $\text{Chi}^2 = 0.74$; $P = 0.69$). Jolly 2007 reported there to be no between-group difference in mortality at 24-months follow up (RR = 1.99, 95% CI 0.50 to 7.88).

Cardiac events

As two studies reported differing cardiac events during exercise programme: Dalal 2007 (CABG, PTCA) and Jolly 2007 (MI, revascularisations at 12 months and 24 months follow up) we were not able to pool data. No significance difference was found in cardiac events between home-based and centre-based settings in either study.

Withdrawals & adherence

Withdrawal from intervention

Although a number of studies reported drop out rates, the reasons for drop out were often unclear so it was therefore not possible to

consistently estimate the number of patients withdrawing from the cardiac rehabilitation programme for each study. However, using the number of completers i.e. number of patients with outcome data at follow up, we found no difference between home based and centre based programmes (fixed effect RR 1.00, 95% CI 0.97 to 1.04; $\text{Chi}^2=11.44$, $\text{df}=10$ ($P=0.32$), $I^2=13\%$, [Analysis 6.1](#)).

Adherence

Nine out of the 12 included studies reported adherence to the cardiac rehabilitation intervention over the duration of the study ([Table 2](#)). However, there was substantial variation in the way in which adherence was defined and measured and some studies reported more than one measure of adherence. Pooling across studies was therefore deemed inappropriate and instead findings are tabulated. Where not reported in the original publications, between-group P-values were calculated by the authors of this report using methods outlined in the Cochrane Handbook ([Higgins 2008](#)). [Table 2](#) summarises the adherence findings for each individual trial. In four out of the remaining seven studies there was no evidence of a significant difference in adherence between home- and centre-based groups although this was not the case in [Arthur 2002](#) and [Marchionni 2003](#) which both showed adherence at $P \leq 0.05$ in favour of home-based cardiac rehabilitation. No other study reported a significantly higher adherence in the centre-based group compared to home.

Costs and healthcare utilisation

Four studies reported costs ([Table 3](#)). Given the difference in currencies and timing of studies is not possible to directly compare the costs across studies. In three ([Carlson 2000](#); [Dalal 2007](#); [Marchionni 2003](#)) of the four studies the healthcare costs associated with home-based cardiac rehabilitation was lower than centre. However, only in Dalal was this lower cost of home CR shown to be statistically different. [Jolly 2007](#) found the costs of home CR to be more expensive than centre-based one although if patient costs were included the costs of the two were the same. Overall healthcare costs were not different between home and centre. Six studies reported different aspects of healthcare resource consumption that included rehospitalisations, primary care consultations and use of secondary care medication ([Table 4](#)). No significant between group differences were seen.

DISCUSSION

The mainstay approach to cardiac rehabilitation delivery in many countries is an inpatient and outpatient hospital-based provision, which often takes place in a supervised University, hospital or community setting. The availability of home-based programmes may provide an opportunity to widen access and participation to

cardiac rehabilitation and thereby may improve uptake and adherence. Home-based cardiac rehabilitation may appear to be a less costly alternative for healthcare economies than more traditional hospital-based. UK figures suggest that 20% of cardiac rehabilitation programmes are currently home-based. This review assessed the randomised controlled trial evidence comparing outcomes of home- and centre-based cardiac rehabilitation.

Our systematic review found no evidence to support a difference in outcomes in cardiac patients receiving home-based or centre-based cardiac rehabilitation either in the short-term (3-12 months) or longer-term (up to 24-months). The study population in the trials were mainly male with a mean age of 51.6 to 69 years. Outcomes considered in this review included exercise capacity, modifiable risk factors (blood pressure, blood lipids and smoking), health-related quality of life, cardiac events (including mortality, revascularisations and rehospitalisations) and adherence. Although some results (diastolic blood pressure, HDL-cholesterol) seemed to have statistical significance, after excluding the most outlining study ([Kassaiyan 2000](#)) the statistically significant difference between groups has been lost. Although not the primary focus of this review, in accord with the two Cochrane reviews of exercise-based cardiac rehabilitation ([Jolliffe 2001](#); [Rees 2004a](#)) we found there to be an improvement in the above following both home- and centre-based cardiac rehabilitation. Healthcare costs appear to depend on the healthcare economy in which cardiac rehabilitation provision is made. However, this review found no consistent evidence to support an important difference in healthcare costs of providing home- versus centre-based programmes.

Our findings are consistent with the previous non-Cochrane systematic review by Jolly ([Jolly 2006](#)). However, this updated review substantially increases the body of evidence base for home-versus centre-based cardiac rehabilitation, incorporating emerging evidence. Our review identified 12 randomised controlled trials in 1,938 cardiac patients, most of them performed during the last eight years, compared with the previous six trials in 749 patients. The Jolly review was critical of the variety of home-based cardiac rehabilitation interventions, the small and poor quality of trials. More recently, two relatively large and high quality UK NHS funded randomised controlled trials comparing home- and hospital-based cardiac rehabilitation have been published ([Dalal 2007](#); [Jolly 2007](#)). The model of home-based provision in the largest three included trials ([Bell 1998](#); [Dalal 2007](#); [Jolly 2007](#)) was the Heart Manual, a home-based cardiac rehabilitation programme that consists of a self-help manual supported by a nurse facilitator ([Lewin 1992](#)).

Our review has limitations. The recruitment of the included trials was primarily limited to stable CHD patients either following an acute-MI or revascularisation. Only one trial was found comparing centre- and home-based cardiac rehabilitation in patients with heart failure. Although the majority of patients in this review were exposed to the Heart Manual model of home-based cardiac

rehabilitation there was evidence of considerable statistically heterogeneity across a number of outcomes across trials. This heterogeneity may well reflect the variety of centre-based cardiac rehabilitation interventions. Trials were pooled using a random effects meta-analysis in the presence of statistical heterogeneity. The majority of studies were of relatively short duration, only one trial reporting outcomes at 24-months (Jolly 2007).

It has been hypothesised that patient preference may have an impact on uptake and adherence to home-based cardiac rehabilitation and there is evidence that white patients who work full- or part-time and who perceive time constraints are more likely to have a preference for home-based provision (Grace 2005). However, such a hypothesis is difficult to test in a traditional RCT designed study and therefore our finding of similar adherence between home and centre needs to be interpreted with caution. The included CHARMS study (Dalal 2007) employed a comprehensive cohort design in addition to the randomised element of home and centre allocation in which there was also a patient preference element (patients could choose between home and hospital-based cardiac rehabilitation). The authors reported outcomes to be very similar between the home and hospital preference arms for all the primary and secondary outcomes to that of the randomised comparison. Adherence to home-based cardiac rehabilitation was also comparable between the randomised (75%) and preference arms (73%). This finding does not support the hypothesis that patients who can choose a programme to suit their lifestyle and preferences will have a higher adherence rate and improved outcomes. As with the randomised comparison, the numbers in the preference arms were small (n=126).

AUTHORS' CONCLUSIONS

Implications for practice

Trials of home- and hospital or centre-based cardiac rehabilitation

have been often been poorly reported and therefore their risk of bias is difficult to ascertain. These trials were mainly conducted on males with a mean age of 52-69 years and may not be generalisable to the wider community of cardiac patients. However, cardiac rehabilitation in both settings appears to be equally effective in improving the clinical and health-related quality of life outcomes in acute MI, revascularisation and heart failure patients. This finding together with an absence of evidence of difference in healthcare costs between the two approaches would support the extension of home-based cardiac rehabilitation programmes such as the Heart Manual. The choice of participating in a more traditional supervised centre-based or home-based programme should reflect the preference of the individual patient, and may have an impact on uptake of cardiac rehabilitation in the individual case.

Implications for research

Data are needed to determine whether the effects of home- and centre-based cardiac rehabilitation reported in short-term trials can be confirmed in the longer term. Further comparative trials are needed to assess the relative impact of supervised centre- versus home-based cardiac rehabilitation in patients with heart failure and chronic angina pectoris. Such studies need to consider economic factors and patient-related outcomes including costs to the healthcare system and health-related quality of life.

ACKNOWLEDGEMENTS

We wish to express our thanks to Dr Philippa Davies who undertook selection of updated titles and abstracts from updated searches, Drs Dalal, Daskapan and Jolly for providing additional details on his included trial and Sue Whiffen for her administrative assistance and the authors of the included studies who provided additional information for the purposes this review.

REFERENCES

References to studies included in this review

Arthur 2002 *{published data only}*

* Arthur HM, Smith KM, Kodis J, McKelvie R. A controlled trial of hospital versus home-based exercise in cardiac patients. *Medicine and Science in Sports and Exercise* 2002;**34**(10):1544–50.

Smith KM, Arthur HM, McKelvie RS, Kodis J. Differences in sustainability of exercise and health-related quality of life outcomes following home or hospital-based cardiac rehabilitation. *European Journal of Cardiovascular Prevention and Rehabilitation* 2004;**11**(4):313–9.

Bell 1998 *{published data only}*

Bell JM. A comparison of a multi-disciplinary home based cardiac rehabilitation programme with comprehensive conventional rehabilitation in post-myocardial infarction patients. Thesis submitted to the University of London for the degree of Doctor of Philosophy 1998.

Carlson 2000 *{published data only}*

Carlson J. A comparison of traditional and modified cardiac rehabilitation protocols on compliance to exercise, patient self-efficacy, cardiovascular outcomes, and program cost. *Dissertation Abstracts International: Section B: The Sciences and Engineering* 1999;**59**(7-B):3388.

* Carlson JJ, Johnson JA, Franklin BA, VanderLaan RL.

- Program participation, exercise adherence, cardiovascular outcomes, and program cost of traditional versus modified cardiac rehabilitation. *American Journal of Cardiology* 2000; **86**(1):17–23.
- Carlson JJ, Norman GJ, Feltz DL, Franklin BA, Johnson JA, Locke S. Self-efficacy, psychosocial factors, and exercise behavior in traditional versus modified cardiac rehabilitation. *Journal of Cardiopulmonary Rehabilitation* 2001; **21**(6):363–73.
- Dalal 2007** *{published and unpublished data}*
- * Dalal HM, Evans PH, Campbell JL, Taylor RS, Watt A, Read KL, et al. Home-based versus hospital-based rehabilitation after myocardial infarction: A randomized trial with preference arms—Cornwall Heart Attack Rehabilitation Management Study (CHARMS). *International Journal of Cardiology* 2007; Vol. 119, issue 2: 202–11.
 - Taylor RS, Watt A, Dalal HM, Evans PH, Campbell JL, Read KL, et al. Home-based cardiac rehabilitation versus hospital-based rehabilitation: a cost effectiveness analysis. *International Journal of Cardiology* 2007; **119**(2):196–201.
- Daskapan 2005** *{published data only}*
- * Daskapan A, Arikan H, Caglar N, Tunalı N, Ataman S. Comparison of supervised exercise training and home-based exercise training in chronic heart failure. *Saudi Medical Journal* 2005; **26**(5):842–7.
 - Daskapan A, Arikan H, Caglar N, Turkman N. The effects of two different exercise programs on pulmonary function tests and quality of life in chronic heart failure: a pilot study (Turkish). *Fizyoterapi Rehabilitasyon* 2005; **16**(2):74–81.
- Gordon 2002 Community** *{published data only}*
- Gordon NF, English CD, Contractor AS, Salmon RD, Leighton RF, Franklin BA, et al. Effectiveness of three models for comprehensive cardiovascular disease risk reduction. *American Journal of Cardiology* 2002; **89**(11):1263–8.
- Gordon 2002 Supervised** *{published data only}*
- Gordon NF, English CD, Contractor AS, Salmon RD, Leighton RF, Franklin BA, et al. Effectiveness of three models for comprehensive cardiovascular disease risk reduction. *American Journal of Cardiology* 2002; **89**(11):1263–8.
- Jolly 2007** *{published and unpublished data}*
- Jolly K, Lip GY, Sandercock J, Greenfield SM, Raftery JP, Mant J, et al. Home-based versus hospital-based cardiac rehabilitation after myocardial infarction or revascularisation: design and rationale of the Birmingham Rehabilitation Uptake Maximisation Study (BRUM): a randomised controlled trial [ISRCTN72884263]. *BMC Cardiovascular Disorders* 2003; **3**:10.
- Jolly K, Lip GY, Taylor RS, Raftery JP, Mant JW, Lane D, et al. The Birmingham Rehabilitation Uptake Maximisation study (BRUM): a randomised controlled trial comparing home-based with centre-based cardiac rehabilitation. *Heart* 2008 Mar 10; **Epub ahead of print**.
- * Jolly K, Taylor R, Lip GY, Greenfield S, Raftery J, Mant J, et al. The Birmingham Rehabilitation Uptake Maximisation Study (BRUM). Home-based compared with hospital-based cardiac rehabilitation in a multi-ethnic population: cost-effectiveness and patient adherence. *Health Technology Assessment* 2007; Vol. 11, issue 35:1–118.
- Kassaian 2000** *{published data only}*
- Kassaian M, Maleki M, Noohi F, Eftekharzadeh M, Arya A, Roshanali F, Momtahan M. Comparing effects of supervised vs. home-based cardiac rehabilitation. *Iranian Heart Journal* 2000; **1**(2):95–102.
- Marchionni 2003** *{published and unpublished data}*
- Marchionni N, Fattirolli F, Fumagalli S, Oldridge N, Del Lungo F, Morosi L, et al. Improved exercise tolerance and quality of life with cardiac rehabilitation of older patients after myocardial infarction: results of a randomized, controlled trial. *Circulation* 2003; Vol. 107, issue 17: 2201–6.
- Miller 1984 Brief** *{published data only}*
- DeBusk RF, Haskell WL, Miller NH, Berra K, Taylor CB, Berger WE III, et al. Medically directed at-home rehabilitation soon after clinically uncomplicated acute myocardial infarction: a new model for patient care. *American Journal of Cardiology* 1985; **55**(4):251–7.
- * Miller NH, Haskell WL, Berra K, DeBusk RF. Home versus group exercise training for increasing functional capacity after myocardial infarction. *Circulation* 1984; **70**(4):645–9.
 - Taylor CB, Houston-Miller N, Ahn DK, Haskell W, DeBusk RF. The effects of exercise training programs on psychosocial improvement in uncomplicated postmyocardial infarction patients. *Journal of Psychosomatic Research* 1986; **30**(5): 581–7.
- Miller 1984 Expanded** *{published data only}*
- DeBusk RF, Haskell WL, Miller NH, Berra K, Taylor CB, Berger WE III, et al. Medically directed at-home rehabilitation soon after clinically uncomplicated acute myocardial infarction: a new model for patient care. *The American Journal of Cardiology* 1985; **55**(4):251–7.
- * Miller NH, Haskell WL, Berra K, DeBusk RF. Home versus group exercise training for increasing functional capacity after myocardial infarction. *Circulation* 1984; **70**(4):645–9.
 - Taylor CB, Houston-Miller N, Ahn DK, Haskell W, DeBusk RF. The effects of exercise training programs on psychosocial improvement in uncomplicated postmyocardial infarction patients. *Journal of Psychosomatic Research* 1986; **30**(5): 581–7.
- Sparks 1993** *{published data only}*
- Sparks KE, Shaw DK, Eddy D, Hanigosky P, Vantrese J. Alternatives for cardiac rehabilitation patients unable to return to a hospital-based program. *Heart Lung* 1993; **22**(4):298–303.
- Wu 2006** *{published data only}*
- Wu SK, Lin YW, Chen CL, Tsai SW. Cardiac rehabilitation vs. home exercise after coronary artery bypass graft surgery: a comparison of heart rate recovery. *American Journal of Physical Medicine and Rehabilitation* 2006; **85**(9):711–7.

References to studies excluded from this review

Ades 2000 *{published data only}*

Ades PA, Pashkow FJ, Fletcher G, Pina IL, Zohman LR, Nestor JR. A controlled trial of cardiac rehabilitation in the home setting using electrocardiographic and voice transtelephonic monitoring. *American Heart Journal* 2000; **139**(3):543–8.

Senuzun 2006 *{published data only}*

Senuzun F, Fadiloglu C, Burke L, Payzin S. Effects of home-based cardiac exercise program on the exercise tolerance, serum lipid values and self-efficacy of coronary patients. *European Journal of Cardiovascular Prevention and Rehabilitation* 2006; **13**(4):640–5.

Sinclair 2005 *{published data only}*

Sinclair A, Conroy S, Davies M, Bayer A. Post-discharge home-based support for older cardiac patients: a randomised controlled trial. *Age and Ageing* 2005; **34**(4):338–43.

Tygesen 2001 *{published data only}*

Tygesen H, Wettervik C, Wennerblom B. Intensive home-based exercise training in cardiac rehabilitation increases exercise capacity and heart rate variability. *International Journal of Cardiology* 2001; **79**(2-3):175–82.

Additional references**Allender 2008**

Allender S, Peto V, Scarborough P, Kaur A, Rayner M. *Coronary heart disease statistics*. London: British Heart Foundation, 2008.

Balady 2007

Balady GJ, Williams MA, Ades PA, Bittner V, Comoss P, Foody JM, et al. Core components of cardiac rehabilitation/secondary prevention programs: 2007 update: a scientific statement from the American Heart Association Exercise, Cardiac Rehabilitation, and Prevention Committee, the Council on Clinical Cardiology; the Councils on Cardiovascular Nursing, Epidemiology and Prevention, and Nutrition, Physical Activity, and Metabolism; and the American Association of Cardiovascular and Pulmonary Rehabilitation. *Circulation* 2007; **115**(20):2675–82.

Bergner 1976

Bergner M, Bobbitt RA, Pollard WE, Martin DP, Gilson BS. The sickness impact profile: validation of a health status measure. *Medical Care* 1976; **14**:57–67.

Beswick 2004

Beswick AD, Rees K, Griebisch I, Taylor FC, Burke M, West RR, et al. Provision, uptake and cost of cardiac rehabilitation programmes: improving services to under-represented groups. *Health Technology Assessment* 2004; **8**(41):1–166.

Davies 2008

Davies P, Beswick A, Wise F, Harris M, Moxham T, Taylor F, Taylor RS. Promoting patient uptake and adherence in cardiac rehabilitation (Protocol). *Cochrane Database of Systematic Reviews* 2008, Issue 2. [DOI: 10.1002/14651858.CD007131]

EuroQoL 1990

The EuroQol Group. EuroQol - a new facility for the measurement of health-related quality of life. *Health Policy* 1990; **16**(3):199–208.

Graham 2007

Graham I, Atar D, Borch-Johnsen K, Boysen G, Burell G, Cifkova R, et al. European guidelines on cardiovascular disease prevention in clinical practice: full text. Fourth Joint Task Force of the European Society of Cardiology and other societies on cardiovascular disease prevention in clinical practice (constituted by representatives of nine societies and by invited experts). *European Journal of Cardiovascular Prevention and Rehabilitation* 2007; **14**(Suppl 2):S1–113.

Heart Manual 2008

Heart Manual Office. *Heart Manual* (<http://www.theheartmanual.com>). Edinburgh: Heart Manual Office, 2008.

Higgins 2008

Higgins JPT, Green S, editors. *Cochrane Handbook for Systematic Reviews of Interventions* 4.2.6 [updated September 2006]. <http://www.cochrane.org/resources/handbook/hbook.htm> [accessed 15 February 2008].

Hunt 1980

Hunt SM, McEwan T. The development of a subjective health indicator. *Sociology of Health and Illness* 1980; **2**:231–246.

Höfer 2004

Höfer S, Lim L, Guyatt G, Oldridge N. The MacNew Heart Disease health-related quality of life instrument: a summary. *Health Quality of Life Outcomes* 2004; **2**:3.

Jolliffe 2001

Jolliffe JA, Rees K, Taylor RS, Thompson D, Oldridge N, Ebrahim S. Exercise-based rehabilitation for coronary heart disease. *Cochrane Database of Systematic Reviews* 2001, Issue 1. [DOI: 10.1002/14651858.CD001800]

Jolly 2006

Jolly K, Taylor RS, Lip GY, Stevens A. Home-based cardiac rehabilitation compared with centre-based rehabilitation and usual care: a systematic review and meta-analysis. *International Journal of Cardiology* 2006; **111**(3):343–51.

Lewin 1992

Lewin B, Robertson IH, Cay EL, Irving JB, Campbell M. Effects of self-help post-myocardial-infarction rehabilitation on psychological adjustment and use of health services. *Lancet* 1992; **339**(8800):1036–40.

McHorney 1993

McHorney CA, Ware JE Jr, Raczek AE. The MOS 36-Item Short-Form Health Survey (SF-36): II. Psychometric and clinical tests of validity in measuring physical and mental health constructs. *Medical Care* 1993; **31**(3):246–63.

NICE 2007

National Institute for Health and Clinical Excellence. MI: Secondary prevention. Secondary prevention in primary and secondary care for patients following a myocardial infarction. Available at <http://www.nice.org.uk/CG48>

[accessed 18 2 2008]. London, UK: National Institute for Health and Clinical Excellence, 2007.

Rees 2004a

Rees K, Taylor RS, Singh S, Coats AJS, Ebrahim S. Exercise-based rehabilitation for heart failure. *Cochrane Database of Systematic Reviews* 2004, Issue 3. [DOI: 10.1002/14651858.CD003331.pub2]

Rees 2004b

Rees K, Bennett P, West R, Davey Smith G, Ebrahim S. Psychological interventions for coronary heart disease. *Cochrane Database of Systematic Reviews* 2004, Issue 2. [DOI: 10.1002/14651858.CD002902.pub2]

Stone 2005

Stone JA, Arthur HM. Canadian guidelines for cardiac rehabilitation and cardiovascular disease prevention, second edition, 2004: executive summary. *Canadian Journal of Cardiology* 2005;**21**(Suppl D):3D–19D.

Taylor 2004

Taylor RS, Brown A, Ebrahim S, Jolliffe J, Noorani H, Rees K, et al. Exercise-based rehabilitation for patients with coronary heart disease: systematic review and meta-analysis of randomized controlled trials. *American Journal of Medicine* 2004;**116**(10):682–92.

WHO 2004

World Health Organization. *The Atlas of Heart Disease and Stroke*. Geneva: World Health Organization, 2004.

References to other published versions of this review

Dalal 2010

Dalal HM, Zawada A, Jolly K, Moxham T, Taylor RS. Home based versus centre based cardiac rehabilitation: Cochrane systematic review and meta-analysis. *BMJ* 2010; **340**:b5631.

* Indicates the major publication for the study

CHARACTERISTICS OF STUDIES

Characteristics of included studies [ordered by study ID]

Arthur 2002

Methods	RCT parallel group	
Participants	n = 120 pts in Home-based CR group; n = 122 pts in Centre-based CR group; 100% post-CABG surgery; mean age 63.3 (SD 13); 81% male <i>Inclusion:</i> 35-49 days post-CABG, able to achieve 40-80% of age/sex-predicted METs on cycle ergometry, read/write English <i>Exclusion:</i> recurrent angina, positive graded exercise test, unable to attend rehabilitation 3x/week, physical limitations, previously participant of out-patient cardiac rehabilitation	
Interventions	<p>Home-based CR (intervention): Exercise: <i>Total duration:</i> 6-months; <i>frequency:</i> 5 sessions/wk; <i>duration:</i> 40 min/session; <i>intensity:</i> 60-70% VO₂max; modality: walking. Also attended 1-hr exercise consultation with exercise specialist at baseline & after 3 month training & completed exercises log which reviewed every 2-months & telephone support call every 2 wks Other: dietary advice & psychological support</p> <p>Centre-based CR (control): Exercise: <i>Total duration:</i> 6-months; <i>frequency:</i> 3 sessions/wk; <i>duration:</i> 40 min/session; <i>intensity:</i> 60-70% VO₂max; modality: cycle ergometer, treadmill, track walking & stair climbing. Supervised by exercise specialist & completed exercises log which reviewed every 1-month Other: dietary advice & psychological support</p>	
Outcomes	<i>Primary:</i> Exercise capacity (METs) <i>Secondary:</i> HRQoL (SF-36); cardiac morbidity, mortality	
Follow up	6- & 18-months post randomisation	
Subgroup analyses	No subgroups described or reported	
Country & settings	Canada, single centre	
Notes		
<i>Risk of bias</i>		
Bias	Authors' judgement	Support for judgement
Adequate sequence generation?	Unclear risk	Not reported
Allocation concealment?	Low risk	"the data analyst, who had no role in this project, prepared the randomization schedule using a blocked format"; "the resulting group assignments were than sealed in opaque envelopes that were opened in sequence after consent"

Arthur 2002 (Continued)

Blinding? All outcomes	Low risk	<i>"the physicians who evaluated the primary variables were blind to the patients assignment"</i>
Incomplete outcome data addressed? All outcomes	Low risk	CONSORT flow diagram shows loss to follow up 20/242 (92%) at 6-months follow up & 24/242 (90%) at 18-months follow up. No imputation of missing data undertaken
Free of selective reporting?	Low risk	All outcomes described in the methods section are reported in the results
Groups balanced at baseline?	High risk	<i>"There were statistically significant differences at baseline between the two groups in weight, resting heart rate, and social support."</i>
Intention to treat analysis?	Low risk	<i>"Analyses were performed based on an intention-to-treat approach."</i>
Groups received same intervention?	Low risk	<i>"Similar numbers of patients in the Hosp and Home groups chose to consult with either clinic dietician or psychologist."</i>

Bell 1998

Methods	RCT parallel group
Participants	n = 152 pts in Home-based CR group; n = 100 pts in Centre-based CR group; 100% acute MI; mean age 59 (SD 8.9); 77% male <i>Inclusion:</i> Acute MI (2 of: elevated serum creatinine kinase or oxaloacetic transaminase, prolonged chest pain consistent with AMI, new Q waves or evolutionary ST changes in ECG) <i>Exclusion:</i> physical infirmity, unable to speak or read English, dementia or psychosis, age >75 y, living >20 miles from CCU, serious persisting medical complications, any other excluding conditions (consultants opinion) (for some hospitals - participation in the previous rehab. programme)
Interventions	Home-based CR (intervention) - Heart Manual: Exercise: Overall duration: 6-weeks; Frequency: not reported; Session duration: not reported; Intensity: not reported Other: 4 phone calls by facilitator, health education, stress management Centre-based CR (control): Exercise: Overall duration & frequency: 12 weeks of 1 session/wk or 4 weeks of 2 sessions/wk; Session duration: ≥20 min; Intensity: 3-4 on Borg RPE scale Other: Education sessions - CHD causes, medication, risk factor modification, stress management & exercise
Outcomes	<i>Primary:</i> Exercise capacity (METs) <i>Secondary:</i> total cholesterol; systolic blood pressure; health-related quality of life (Nottingham Health Profile, NHP); smoking Mortality; readmission rate; use of primary care services

Bell 1998 (Continued)

Follow up	16 & 48 weeks post randomisation (20 & 52 weeks post MI)	
Subgroup analyses	No subgroups described or reported	
Country & settings	UK, 5 district hospitals	
Notes	Published as PhD thesis only	
<i>Risk of bias</i>		
Bias	Authors' judgement	Support for judgement
Adequate sequence generation?	Unclear risk	Not reported
Allocation concealment?	Low risk	<i>"Series of sealed envelopes containing cards evenly distributed between conditions ... envelopes were taken sequentially ... opened envelopes were retained and returned to trial coordinator"</i>
Blinding? All outcomes	Low risk	<i>"All measurements were performed 'blind' by members of the medical staff and technicians"</i>
Incomplete outcome data addressed? All outcomes	Unclear risk	Follow up data on all randomised patient is not reported. No CONSORT flow diagram is reported and it is difficult to determine from report those were loss to follow up or dropped out
Free of selective reporting?	Low risk	All outcomes described in the methods section are reported in the results
Groups balanced at baseline?	Low risk	There were no statistically significant differences in population demographics between two groups
Intention to treat analysis?	High risk	<i>"Statistical analysis followed the intention-to-treat principle ... on the basis 7 subjects randomised to conventional treatment [centre-based CR] were classified as non-compliers"</i> and excluded from analysis
Groups received same intervention?	High risk	Although the intervention to both groups consisted of exercise, education and stress management, the nature and amount of intervention was quite different

Carlson 2000

Methods	RCT parallel group
Participants	n = 38 pts in Home-based CR group; n = 42 pts in Centre-based CR group; diagnosis: coronary artery bypass, angioplasty, MI, angiographically confirmed CHD; case mix: MI, revascularisation & CHD; mean age 59 (SD 14); 83% male

	<p><i>Inclusion:</i> men & women 35-75yrs referred for first time to outpatient cardiac rehabilitation, living ≤30 miles from the rehab. facility, of low-to-moderate cardiac risk <i>Exclusion:</i> not reported</p>	
Interventions	<p>Home-based CR (intervention): Exercise: Overall duration: 25 weeks; Frequency: 2-5 sessions/wk; Duration: 30-40 mins/session; Intensity: 60-85% aerobic capacity; Modality: aerobic exercise. First 4 wks - 3 hospital based exercise session/week with ECG monitoring & then progressively reduce frequency of centre-based sessions Other: weekly educational & counselling meetings that included sessions on exercise, diet, risk factor, drugs and over coming barriers to behaviour change. Based on Bandura's self-efficacy theory Centre-based CR (control): Exercise: Overall duration: 25 weeks; Frequency: 2-3 sessions/wk; Duration: 30-45 min/session; Intensity: 60-85% aerobic capacity; Modality: aerobic exercise Other: 3 sessions of education & counselling that included sessions on exercise, diet, risk factor & drugs</p>	
Outcomes	<p><i>Primary:</i> Peak functional capacity (METs), LDL-cholesterol <i>Secondary:</i> Total cholesterol, HDL-cholesterol, triglycerides, blood pressure, cardiovascular medications, costs, adherence (exercise sessions attended)</p>	
Follow up	6-months post randomisation	
Subgroup analyses	No subgroups described or reported	
Country & settings	USA, single hospital centre	
Notes		
Risk of bias		
Bias	Authors' judgement	Support for judgement
Adequate sequence generation?	Unclear risk	Not reported
Allocation concealment?	Unclear risk	Not reported
Blinding? All outcomes	Unclear risk	"it was not possible to blind the clinicians to the protocol patients were assigned" Outcome blinding not reported
Incomplete outcome data addressed? All outcomes	High risk	"significantly more TP [centre-based CR] participants dropped out" Because of more TP [centre-based CR] participants dropped out and failed to return for their 6-month [exercise test] evaluation, this evaluation is a representation of more compliant patients"

Carlson 2000 (Continued)

Free of selective reporting?	Low risk	All outcomes described in the methods section are reported in the results
Groups balanced at baseline?	Low risk	“...only significant difference between groups was a higher resting systolic blood pressure in the former [centre-based CR] ...selected demographic and psychological measures including socioeconomic status and social support were comparable between the 2 groups at baseline”
Intention to treat analysis?	Low risk	The conduction of intention to treat analysis not stated directly, but results appear to be presented according to original random allocation
Groups received same intervention?	High risk	“The primary differences in the MP [home-based CR] compared with the TP [centre-based CR] included: ... (2) an ongoing weekly education/support group, and (3) education and counselling that emphasized overcoming barriers associated with developing independent exercise and nutrition behaviours” Although both groups both received exercise training, education and counselling the amount and nature of this intervention was different between groups

Dalal 2007

Methods	RCT parallel group
Participants	n = 60 pts in Home-based CR group; n = 44 pts in Centre-based CR group; 100% post-MI; mean age 62 (SD 15); 81% male <i>Inclusion:</i> Confirmed acute myocardial infarction (WHO criteria), ability to read English, registered with GP in one of two primary care trusts <i>Exclusion:</i> Severe heart failure, unstable angina, uncontrolled arrhythmia, history of major psychiatric illness, other significant co-morbidity precluding the ability to exercise on the treadmill, patients readmitted with acute myocardial infarction who had already received an intervention earlier in the study
Interventions	Home-based CR (intervention) - Heart Manual: Exercise: Overall duration: 6-wks; frequency: not reported; duration: not reported; intensity: not reported; modality: walking. Home visit in 1 st week after discharge by CR nurse followed up by up to 4 telephone calls at 2, 3, 4, & 6 wks Other: Stress management & education Centre-based CR (control): Exercise: Overall duration: 8-10 wks; frequency: 1-5 sessions/wk; duration: not reported; modality: not reported. Supervised and group based Other: Input from dietician, psychologist, occupational therapist & pharmacist
Outcomes	<i>Primary:</i> Quality of life (MacNew questionnaire), total cholesterol <i>Secondary:</i> exercise capacity (METs), self-reported smoking Cardiovascular morbidity, mortality, secondary prevention medication use

Dalal 2007 (Continued)

Follow up	9-months post randomisation	
Subgroup analyses	No subgroup analyses described or reported	
Country & settings	UK, single centre	
Notes		
Risk of bias		
Bias	Authors' judgement	Support for judgement
Adequate sequence generation?	Low risk	"computerised random number trial allocation sequence was determined before the study"
Allocation concealment?	Low risk	"allocation was transferred to sequentially numbered, opaque, sealed envelopes and concealed from the research nurse, who carried out baseline assessment"
Blinding? All outcomes	Low risk	"the person assessing the primary outcome questionnaires was blinded to allocation"
Incomplete outcome data addressed? All outcomes	Low risk	"the last known observation carried forward to replace missing values at 9 months for the primary outcome measures."
Free of selective reporting?	Low risk	All outcomes described in the methods section are reported in the results
Groups balanced at baseline?	Low risk	"The randomized groups were well balanced, apart from a higher proportion of patients in employment in the home based group (51% versus 26%, $p=0.013$)"
Intention to treat analysis?	Low risk	"Data were analyzed on an intention to treat principle"
Groups received same intervention?	Low risk	

Daskapan 2005

Methods	RCT parallel group
Participants	n = 15 patients in Home-based CR group; n = 14 pts in Centre-based CR group; diagnosis: heart failure; case mix: CHF class II or III NYHA with ischemic or idiopathic dilated cardiomyopathy; mean age 52 (SD 8.5); 76% male <i>Inclusion:</i> heart failure of >3 month duration <i>Exclusion:</i> valvular heart disease, exercise-induced cardiac arrhythmias, symptomatic myocardial ischemia within 3 months, taking beta-blockers

Interventions	<p>Home-based CR (intervention): Exercise: Total duration: 12 weeks; frequency: 3 sessions/wk; duration: 45 mins/session (including warm-up, cool-down, recovery); intensity: up to 60% peak heart rate (RPE 12-16); modality: walking; follow-up logs completed daily/returned biweekly. Weekly phone calls from staff monitoring adherence & progress, monthly phone calls from patients for control purposes Other: not reported Centre-based CR (control): Exercise: Total duration: 12 weeks; frequency: 3 sessions/wk; duration: 45 mins/session (including warm-up, cool-down, recovery); intensity: 60% peak heart rate; modality: walking on a treadmill. Supervised Other: not reported</p>	
Outcomes	(primary and secondary outcomes not distinguished) exercise capacity (ml/kg/min), resting BP systolic & diastolic BP, adherence, dropouts; additionally data on mortality obtained by personal contact	
Follow up	12 weeks post randomisation	
Subgroup analyses	No subgroups described or reported	
Country & settings	Turkey, single centre	
Notes	Additional data on mortality obtained by personal contact	
Risk of bias		
Bias	Authors' judgement	Support for judgement
Adequate sequence generation?	Unclear risk	Not reported
Allocation concealment?	Unclear risk	Not reported
Blinding? All outcomes	Unclear risk	Not reported
Incomplete outcome data addressed? All outcomes	Low risk	
Free of selective reporting?	Low risk	All outcomes described in the methods section are reported in the results
Groups balanced at baseline?	Low risk	"Among patients who completed the study, no differences in demographic characteristics were seen between the 2 study groups after randomization ($p>0.05$)."
Intention to treat analysis?	High risk	ITT not reported

Groups received same intervention?	Low risk	“We chose lower intensity ...training prescriptions in the HETG to avoid any adverse occurrences and also in the SETG to provide comparable training intensity levels between 2 groups.”
------------------------------------	----------	--

Gordon 2002 Community

Methods	RCT parallel group
Participants	<p>n = 54 pts in physician-supervised Home-based CR group; n = 49 pts in community Home-based CR group; n = 52 pts in Centre-based CR group; diagnosis: 100% CAD (MI and/or CABG and/or PTCA and/or chronic stable angina); age: 60.4 (SD 9.4); 75% male</p> <p><i>Inclusion:</i> diagnosed CAD (as above); low-moderate risk of cardiac events (1. no cardiac arrest within 1 year, 2. no complex ventricular dysrhythmia, 3. ejection fraction <40%, 4. no complicated MI or cardiac surgery, 5. no increasing systolic BP response to exercise testing, 6. no angina pectoris <5.0 METs); ≥4 weeks posthospitalization; age 21-75 y; no life-threatening illness and/or psychological abnormality; speak/write English; ability to complete exercise treadmill test; ability to attend 36 cardiac rehabilitation sessions</p> <p><i>Exclusion:</i> not defined</p>
Interventions	<p>Home-based CR</p> <p>Group I (Supervised Home-based CR)</p> <p>Exercise: Total duration: 12 weeks; frequency & intensity: individually prescribed (30-60 min of aerobic exe, 60-85% peak HR), gradually updated; appointments: 2 office visits, 4 phone calls</p> <p>Other: written materials, audiotapes, nutrition, weight & stress management, smoking cessation programme, individual CAD risk factors management</p> <p>Group II (Community Home-based CR)</p> <p>Exercise: Total duration: 12 weeks; frequency & intensity: individually prescribed (30-60 min of aerobic exe, 60-85% peak HR), gradually updated; appointments: 12 on site visits or telephone calls (patient choice)</p> <p>Other: written materials, audiotapes, nutrition, weight & stress management, smoking cessation programme, individual CAD risk factors management</p> <p>Centre-based CR</p> <p>Exercise: Total duration: 12 weeks; frequency: 3 sessions/wk (total of 36 sessions = appointments); intensity: individually prescribed (30-60 min of aerobic exe, 60-85% peak HR); continuous ECG telemetry during exercise</p> <p>Other: written materials, audiotapes, education on CAD risk factors & lifestyle modification</p>
Outcomes	(primary & secondary risk factors not distinguished) maximal oxygen uptake, blood pressure, fasting serum lipids, self-reported smoking status, rehospitalization, adherence (completion of appointments)
Follow up	12 weeks post randomisation
Subgroup analyses	Changes reported for all patients and for patients with baseline values defined as abnormal

Gordon 2002 Community (Continued)

Country & settings	USA single centre	
Notes		
<i>Risk of bias</i>		
Bias	Authors' judgement	Support for judgement
Adequate sequence generation?	Unclear risk	Not reported
Allocation concealment?	Unclear risk	Not reported
Blinding? All outcomes	Unclear risk	Not reported
Incomplete outcome data addressed? All outcomes	Low risk	Data for 142 pts who completed exercise testing at baseline and at follow up (not all 155 pts randomised) reported only; numbers of dropouts reported and reasons described
Free of selective reporting?	Low risk	All outcomes mentioned in methods are reported in results
Groups balanced at baseline?	Low risk	<i>"Randomization did not result in statistical significant differences among patients assigned to the 3 interventions."</i>
Intention to treat analysis?	Unclear risk	Not reported
Groups received same intervention?	Low risk	

Gordon 2002 Supervised

Methods	see Gordon2002-Commun
Participants	
Interventions	
Outcomes	
Follow up	
Subgroup analyses	
Country & settings	
Notes	
<i>Risk of bias</i>	

Gordon 2002 Supervised (Continued)

Bias	Authors' judgement	Support for judgement
Adequate sequence generation?	Unclear risk	Not reported
Allocation concealment?	Unclear risk	Not reported
Blinding? All outcomes	Unclear risk	Not reported
Incomplete outcome data addressed? All outcomes	Low risk	Data for 142 pts who completed exercise testing at baseline and at follow up (not all 155 pts randomised) reported only; numbers of dropouts reported and reasons described
Free of selective reporting?	Low risk	All outcomes mentioned in methods are reported in results
Groups balanced at baseline?	Low risk	<i>"Randomization did not result in statistical significant differences among patients assigned to the 3 interventions."</i>
Intention to treat analysis?	Unclear risk	Not reported
Groups received same intervention?	Low risk	

Jolly 2007

Methods	RCT parallel group
Participants	n = 263 pts in Home-based CR group; n = 262 pts in Centre-based CR group; diagnosis: MI 256 pts; PTCA 211 pts; CABG 56 pts; mean age 61 (SD 10.8); 77% male; 80.2% white <i>Inclusion:</i> an acute MI, coronary angioplasty (±stenting) or CABG <i>Exclusion:</i> inability to speak either English or Punjabi, dementia, severe hearing impairment, sight defects of sufficient severity to prevent them from reading the Heart Manual & serious persisting complications
Interventions	Home-based CR (intervention) - Heart Manual: Exercise: Overall duration: 6 wks Heart Manual programme & 12 wks nurse support; frequency: up to daily; duration: not reported; intensity: not reported; modality: walking Other: education on risk factors, lifestyle changes, medications & stress management (relaxation tapes) Centre-based CR (control): Exercise: Total duration: 6-12 wks; Frequency: 1 or 2 sessions/wk; duration: 25-30mins/session; intensity: 65-75% HRmax; modality: circuit training, cycle ergometer Other: education & stress management (relaxation)

Outcomes	<p><i>Primary:</i> Serum cholesterol, total, HDL-cholesterol, blood pressure, exercise capacity (incremental shuttle walking test, ISWT), smoking cotinine-validated)</p> <p><i>Secondary:</i> quality of life (EQ-5D), health service utilisation (hospital readmissions, primary care visits, medication)</p> <p>Mortality, cardiovascular events, costs</p>	
Follow up	6, 12 & 24 months	
Subgroup analyses	<p>Yes. <i>“Interaction terms between these factors [diagnosis (MI/revascularisation), age, sex and ethnicity] and rehabilitation setting were included to investigate possible differences in treatment effect between subgroups of patients.”</i></p>	
Country & settings	UK, 4 hospital centres	
Notes		
Risk of bias		
Bias	Authors’ judgement	Support for judgement
Adequate sequence generation?	Low risk	<i>“Patients who consented to randomisation were randomised on an individual basis with minimisation by (1) original diagnosis (MI/revascularisation), (2) age (<50/50-74/75+ years), (3) sex, (4) ethnicity (Caucasian/Asian/other) and (5) hospital of recruitment.”</i>
Allocation concealment?	Low risk	<i>“Allocation was undertaken by the Birmingham Cancer Clinical Trials Unit, a group that was independent from the trial team ... When a patient agreed to be randomised, ... the research nurse telephoned the Clinical Trials Unit, ... and was given an allocation group.”</i>
Blinding? All outcomes	Low risk	<i>“Assessments were blinded, with follow-up undertaken by a research nurse who had neither recruited the patient nor provided home CR support.”</i>
Incomplete outcome data addressed? All outcomes	Low risk	<i>“A sensitivity analysis was undertaken on the 12-month data to assess the potential impact of the missing values for the ISWT, SBP, DBP, TC and the HADS scores.”</i>
Free of selective reporting?	Low risk	All outcomes described in the methods section are reported in the results
Groups balanced at baseline?	Low risk	<i>“Demographic characteristics, diagnosis, past medical history and cardiac risk factors were well matched between the two arms at baseline.”</i>
Intention to treat analysis?	Low risk	<i>“All data were analysed by intention-to-treat (ITT).”</i>

Groups received same intervention?	High risk	Although both groups received exercise, education and stress management, the nature and amount of intervention between groups was different
------------------------------------	-----------	---

Kassaian 2000

Methods	RCT parallel group	
Participants	n = 65 pts in Centre-based CR group (active intervention); n = 60 pts in Home-based CR group (controls); diagnosis: MI 23.2%; CABG 76.8%; case mix: MI: Hosp 32.3%, Home 13.3%; CABG: Hosp 67.7%, Home 86.7%; mean age 55 (SD 9.5); 100% male <i>Inclusion:</i> AMI or CABG in last 1-2 month, NYHA<IV, EF≥30%, able to exercise on a treadmill & participate in exercise programme <i>Exclusion:</i> high-risk stress test, decompensated CHF (NYHA IV), unstable angina, uncontrolled AF, high-grade AV block (grade 2 or 3), active pericarditis or myocarditis, recent pulmonary thromboembolism, exercise-induced asthma, claudication, fixed-rate permanent pacemaker, severe medical problem	
Interventions	<p>Centre-based CR (intervention): Exercise: <i>Total duration:</i> 12 weeks; <i>frequency:</i> 3 sessions/wk; <i>duration:</i> 20-30 min+10 min warm-up+10 min cool-down/session; <i>intensity:</i> 60-85% (not reported if relative to HRmax or V02max); <i>modality:</i> treadmill Other: not reported</p> <p>Home-based CR (control): Exercise: <i>Total duration:</i> 12 weeks; ; <i>frequency:</i> not reported; <i>duration:</i> not reported; <i>intensity:</i> “intensity based on exercise test results” Other: patients have been taught to count their pulse rate</p>	
Outcomes	(primary and secondary outcomes not distinguished) systolic BP, diastolic BP, heart rate (all resting & sub-maximal), functional capacity (METs), BMI, cholesterol: total, LDL, HDL, triglyceride	
Follow up	12 weeks post randomization	
Subgroup analyses	Subgroup analysis: comparison of functional capacity, sub-maximal SBP, DBP & heart rate in pts with left ventricular dysfunction vs good LV function	
Country & settings	Iran, single centre	
Notes		
Risk of bias		
Bias	Authors’ judgement	Support for judgement
Adequate sequence generation?	Unclear risk	Not reported

Kassaian 2000 (Continued)

Allocation concealment?	Unclear risk	Not reported
Blinding? All outcomes	Unclear risk	Not reported
Incomplete outcome data addressed? All outcomes	Unclear risk	No information on loss to follow up, nor on missing data management
Free of selective reporting?	Unclear risk	Not all outcomes reported mentioned in methods section
Groups balanced at baseline?	Low risk	“Among patients who completed the study no differences in demographic characteristics were seen between the two study groups after randomisation.”
Intention to treat analysis?	Unclear risk	Not reported
Groups received same intervention?	Unclear risk	Details of home-based intervention not reported

Marchionni 2003

Methods	RCT parallel group
Participants	n = 90 pts in Home-based CR group; n = 90 pts in Centre-based CR group; 100% MI; mean age 69 (SD 1.6); 71% male <i>Inclusion:</i> age >45 y, MI <i>Exclusion:</i> Severe cognitive impairment; physical disability; left ventricular ejection fraction <35%; contraindications to vigorous exercise; eligibility for myocardial revascularisation, living too far from CR unit
Interventions	Home-based CR (intervention): Exercise: Overall duration; 8-wks; Frequency: 3 days/wk; Duration: 1-hour; Intensity 70-85% peak HR; Modality: cycle ergometer. Physical therapist home visits every other week Other: monthly family-oriented support groups Centre-based CR (control): Exercise: Overall duration; 12-week programme; Frequency: 3 days/wk; Duration: not reported; Intensity 70-85% peak HR; Modality: cycle ergometer. Trans-telephonic ECG monitoring during exercise Other: risk factor management counselling; support group meetings
Outcomes	<i>Primary:</i> Total work capacity (TWC) <i>Secondary:</i> Health-related quality of life (Sickness Impact Profile, SIP), mortality, morbidity (cardiovascular events), healthcare utilisation (medical visits, rehospitalisations), costs & adherence (number of completed training sessions)
Follow up	2, 8 & 14 months post randomisation

Subgroup analyses	Subgroup analysis in age groups (middle-aged 45-65 years, old 65-75 years, very old >75 years)	
Country & settings	Italy, single hospital centre	
Notes	Data presented separately for 3 age groups. Follow up data on charts only; authors contacted for numerical data at follow up & these have been supplied for TWC & SIP separately for 3 groups; data pooled across age groups by reviewers	
<i>Risk of bias</i>		
Bias	Authors' judgement	Support for judgement
Adequate sequence generation?	Unclear risk	Not reported
Allocation concealment?	Unclear risk	Not reported
Blinding? All outcomes	Low risk	"Testing personnel were blinded to patient assignment."
Incomplete outcome data addressed? All outcomes	Low risk	"we performed a sensitivity analysis comparing results obtained with and without replacement of missing data with data obtained with the expectation-maximization imputation method. Because the 2 analyses provided similar results, which were also similar with missing data substituted with data estimated in a worst-case scenario, only the data from patients who completed the study are presented"
Free of selective reporting?	Low risk	All outcomes described in the methods section are reported in the results
Groups balanced at baseline?	Low risk	"baseline sociodemographic and clinical characteristics were similar across the 3 arms of the trial" Baseline characteristics by home and hospital group allocation not reported in tabular format
Intention to treat analysis?	Low risk	Although term 'intention to treat analysis' was not stated directly, on the basis of CONSORT diagram presented, the groups did appear to be analysed according to original random allocation
Groups received same intervention?	Low risk	

Miller 1984 Brief

Methods	RCT parallel group
Participants	n = 66 patients in Home-based CR group (33 in brief exercise programme subgroup & 33 in extended subgroup); n = 61 patients in Centre-based CR group (31 in brief subgroup & 30 in extended subgroup); 100% uncomplicated acute MI; mean age 52 (SD 9); 100% male <i>Inclusion:</i> Uncomplicated AMI (elevated serum creatinine kinase or oxaloacetic transaminase, prolonged chest pain consistent with AMI, new Q waves or evolutionary ST changes in ECG) <i>Exclusion:</i> Unable to undertake exercise test, congestive heart failure, unstable angina pectoris, valvular heart disease, atrial fibrillation, bundle branch block, history of bypass, stroke, orthopaedic abnormalities, peripheral vascular disease, chronic pulmonary obstructive disease, obesity
Interventions	Home-based CR (intervention): Exercise: <i>Overall duration:</i> 8 wks (brief) or 23 wks (extended), <i>Frequency:</i> 5 sessions/wk; <i>Duration:</i> 30 mins/session; <i>Intensity:</i> 70-85% HRmax; <i>modality:</i> stationary cycling. Portable heart rate monitors & teletransmissions of ECG; 2 phone calls/wk by staff to verify training intensity, clinical status and medication Other: No education or psychological intervention reported Centre-based CR (control): Exercise: <i>Overall duration:</i> 8 wks (brief) or 23 wks (extended), <i>Frequency:</i> 5 sessions/wk; <i>Duration:</i> 60 mins/session; <i>Intensity:</i> 70-85% HRmax; <i>modality:</i> walking/jogging. Group based & supervised Other: No education or psychological intervention reported
Outcomes	Exercise capacity; mortality & cardiovascular morbidity
Follow up	23 weeks post randomisation
Subgroup analyses	Yes. Results reported according to the two subgroups reported i.e. brief vs extended exercise training
Country & settings	USA, single hospital centre
Notes	Results of two subgroups included into analysis separately
Risk of bias	
Bias	Authors' judgement Support for judgement
Adequate sequence generation?	Unclear risk Not reported
Allocation concealment?	Unclear risk Not reported
Blinding? All outcomes	Unclear risk Not reported

Miller 1984 Brief (Continued)

Incomplete outcome data addressed? All outcomes	Low risk	Drop out reported. No imputation of missing data discussed. Effect of incomplete outcome data likely to be small
Free of selective reporting?	Low risk	All outcomes described in the methods section are reported in the results
Groups balanced at baseline?	Unclear risk	Baseline characteristics not reported
Intention to treat analysis?	Low risk	Term 'intention to treat analysis' not stated directly, but results appear to be presented according to original random allocation. This is further supported by a trial flow diagram in the paper
Groups received same intervention?	Low risk	Both home and centre groups were very closely balanced in terms of the exercise training received

Miller 1984 Expanded

Methods	see Miller 1984 Brief	
Participants		
Interventions		
Outcomes		
Follow up		
Subgroup analyses		
Country & settings		
Notes		
<i>Risk of bias</i>		
Bias	Authors' judgement	Support for judgement
Adequate sequence generation?	Unclear risk	Not reported
Allocation concealment?	Unclear risk	Not reported
Blinding? All outcomes	Unclear risk	Not reported
Incomplete outcome data addressed? All outcomes	Low risk	Drop out reported. No imputation of missing data discussed. Effect of incomplete outcome data likely to be small

Miller 1984 Expanded (Continued)

Free of selective reporting?	Low risk	All outcomes described in the methods section are reported in the results
Groups balanced at baseline?	Unclear risk	Baseline characteristics not reported
Intention to treat analysis?	Low risk	Term 'intention to treat analysis' not stated directly, but results appear to be presented according to original random allocation. This is further supported by a trial flow diagram in the paper
Groups received same intervention?	Low risk	Both home and centre groups were very closely balanced in terms of the exercise training received

Sparks 1993

Methods	RCT parallel groups	
Participants	n = 10 pts in Home-based CR group; n = 10 pts in Centre-based CR group; diagnosis: MI, CABG, PTCA; case mix: MI & revascularisation; mean age 51.6 (SD 12); 100% male <i>Inclusion:</i> being male cardiac patient <i>Exclusion:</i> not capable for exercising on a bicycle ergometer, serious arrhythmias, symptoms of frequent chest pain, shortness of breath, hypertension	
Interventions	<p>Home-based CR (intervention): Exercise: <i>Overall duration:</i> 12-week programme; <i>Frequency:</i> 3 days/wk; <i>Duration:</i> 1-hour; <i>Intensity</i> 60-75% peak HR; <i>Modality:</i> cycle ergometer. Trans-telephonic ECG monitoring Other: Education materials on diet, medications, risks and benefits of the exercise Centre-based CR (control): As above but no trans-telephonic ECG monitoring during exercise</p>	
Outcomes	Exercise capacity (peak VO ₂ max); adherence (compliance with exercise); safety (drop out)	
Follow up	12 weeks post randomisation	
Subgroup analyses	No subgroups described or reported	
Country & settings	USA, single hospital centre	
Notes	Data read from graphs	
Risk of bias		
Bias	Authors' judgement	Support for judgement
Adequate sequence generation?	Unclear risk	Not reported

Sparks 1993 (Continued)

Allocation concealment?	Unclear risk	Not reported
Blinding? All outcomes	Unclear risk	Not reported
Incomplete outcome data addressed? All outcomes	Low risk	Only 1/20 (5%) drop out reported
Free of selective reporting?	Low risk	All outcomes described in the methods section are reported in the results
Groups balanced at baseline?	Low risk	Although no statement of similarity of baseline characteristics, the characteristic of both groups in table 1 appeared similar
Intention to treat analysis?	Low risk	The conduction of intention to treat analysis not stated directly, but results appear to be presented according to original random allocation
Groups received same intervention?	Low risk	

Wu 2006

Methods	RCT parallel group
Participants	n = 18 pts in Home-based CR group; n = 18 pts in Centre-based CR group; 100% post CABG; mean age 61.9 (SD 7.3); 100% male <i>Inclusion:</i> No previous CABG, no neurologic impairment like stroke/brain injury, no severe musculoskeletal disease, no complications during hospitalisations like infection, shock, arrhythmia, prolonged ventilation <i>Exclusion:</i> uncontrolled dysrhythmia or continuous ventricular tachycardia during exercise testing, no possibility of completing test at discharge or 12 wks later
Interventions	Home-based CR (intervention): Exercise: Total duration: 12 weeks; frequency: ≥ 3 sessions/wk; duration: 30-60 min+10 min warm-up+10 min cool-down/session; intensity: 60-85% HRmax; modality: fast walking or jogging. Exercise documented in record book. Prescription of exercise individually given & updated every 2 wks by rehabilitation nurse Other: not reported Centre-based CR (control): Exercise: Total duration: 12 weeks; frequency: 3 sessions/wk (total 36 sessions); duration: 30-60 min+10 min warm-up+10 min cool-down/session; intensity: 60-85% HRmax; modality: cycle ergometer, treadmill. Exercise supervised by cardiopulmonary physical therapist Other: not reported
Outcomes	(primary and secondary outcomes not distinguished) exercise capacity (METs)
Follow up	12 weeks post randomisation

Subgroup analyses	No subgroups described or reported	
Country & settings	Taiwan (China), single centre	
Notes		
<i>Risk of bias</i>		
Bias	Authors' judgement	Support for judgement
Adequate sequence generation?	Unclear risk	<i>"Subjects were randomly assigned by drawing lots"</i>
Allocation concealment?	Unclear risk	Not reported
Blinding? All outcomes	Low risk	<i>"The evaluators of the exercise stress test were also masked to the group assignments."</i>
Incomplete outcome data addressed? All outcomes	Unclear risk	Not reported
Free of selective reporting?	Low risk	All outcomes described in methods section were reported in results
Groups balanced at baseline?	Low risk	<i>"Randomization did not result in statistical significances among subjects assigned to the three groups."</i>
Intention to treat analysis?	Unclear risk	Not reported, although it appears that patients were analysed according to original allocation
Groups received same intervention?	Low risk	

AMI=acute myocardial infarction
 CABG=coronary artery bypass graft
 CAD=coronary artery disease
 CCU=coronary care unit
 CHD=coronary heart disease
 CHF=congestive heart failure
 CR=cardiac rehabilitation
 ECG=electrocardiogram
 HRQoL=health related quality of life
 ISWT=incremental shuttle walking test
 ITT=intention to treat
 METs=metabolic equivalents
 MI=myocardial infarction
 NYHA=New York Heart Association classification
 PTCA=percutaneous transluminal coronary angioplasty
 RCT=randomised controlled trial

SD=standard deviation

Characteristics of excluded studies *[ordered by study ID]*

Study	Reason for exclusion
Ades 2000	Non RCT
Senuzun 2006	Trial experimental arm received home-based cardiac rehabilitation; the programme issued in control arm was not described
Sinclair 2005	Trial experimental arm received home-based cardiac rehabilitation, while the control group did not receive centre based CR (only 6% [n=12] of the participants in the control group were referred to CR and only 3% [n=8] were known to have attended)
Tygesen 2001	Both trial arms received home-based cardiac rehabilitation

DATA AND ANALYSES

Comparison 1. Exercise capacity

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Exercise capacity 3-12 month	14	1557	Std. Mean Difference (IV, Random, 95% CI)	-0.11 [-0.35, 0.13]
2 Exercise capacity 12-24 month	3	1074	Std. Mean Difference (IV, Fixed, 95% CI)	0.11 [-0.01, 0.23]

Comparison 2. Blood Pressure [mm Hg]

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Systolic BP 3-12 month	8	1053	Mean Difference (IV, Random, 95% CI)	0.58 [-3.29, 4.44]
2 Diastolic BP 3-12 month	7	927	Mean Difference (IV, Fixed, 95% CI)	-1.85 [-2.96, -0.74]

Comparison 3. Blood lipids [mmol/l]

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Total cholesterol 3-12 month	7	1019	Mean Difference (IV, Random, 95% CI)	-0.13 [-0.31, 0.05]
2 HDL cholesterol 3-12 month	5	793	Mean Difference (IV, Fixed, 95% CI)	-0.06 [-0.11, -0.02]
3 LDL-cholesterol 3-12 month	4	324	Mean Difference (IV, Fixed, 95% CI)	-0.15 [-0.31, 0.01]
4 Triglycerides 3-12 month	4	328	Mean Difference (IV, Random, 95% CI)	-0.15 [-0.41, 0.11]

Comparison 4. Smoking

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Smoking 3-12 month	5	922	Risk Ratio (M-H, Random, 95% CI)	1.00 [0.71, 1.41]

Comparison 5. Mortality

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Mortality	4	909	Risk Ratio (M-H, Fixed, 95% CI)	1.31 [0.65, 2.66]

Comparison 6. Completers


Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Completers	13	1620	Risk Ratio (M-H, Fixed, 95% CI)	1.02 [0.99, 1.06]

Analysis 1.1. Comparison 1 Exercise capacity, Outcome 1 Exercise capacity 3-12 month.


Review: Home-based versus centre-based cardiac rehabilitation

Comparison: 1 Exercise capacity

Outcome: 1 Exercise capacity 3-12 month



(... Continued)




Analysis 1.2. Comparison 1 Exercise capacity, Outcome 2 Exercise capacity 12-24 month.

Review: Home-based versus centre-based cardiac rehabilitation

Comparison: 1 Exercise capacity

Outcome: 2 Exercise capacity 12-24 month




Analysis 2.1. Comparison 2 Blood Pressure [mm Hg], Outcome 1 Systolic BP 3-12 month.

Review: Home-based versus centre-based cardiac rehabilitation

Comparison: 2 Blood Pressure [mm Hg]

Outcome: 1 Systolic BP 3-12 month




Analysis 2.2. Comparison 2 Blood Pressure [mm Hg], Outcome 2 Diastolic BP 3-12 month.

Review: Home-based versus centre-based cardiac rehabilitation

Comparison: 2 Blood Pressure [mm Hg]

Outcome: 2 Diastolic BP 3-12 month




Analysis 3.1. Comparison 3 Blood lipids [mmol/l], Outcome 1 Total cholesterol 3-12 month.

Review: Home-based versus centre-based cardiac rehabilitation

Comparison: 3 Blood lipids [mmol/l]

Outcome: 1 Total cholesterol 3-12 month




Analysis 3.2. Comparison 3 Blood lipids [mmol/l], Outcome 2 HDL cholesterol 3-12 month.

Review: Home-based versus centre-based cardiac rehabilitation

Comparison: 3 Blood lipids [mmol/l]

Outcome: 2 HDL cholesterol 3-12 month




Analysis 3.3. Comparison 3 Blood lipids [mmol/l], Outcome 3 LDL-cholesterol 3-12 month.

Review: Home-based versus centre-based cardiac rehabilitation

Comparison: 3 Blood lipids [mmol/l]

Outcome: 3 LDL-cholesterol 3-12 month




Analysis 3.4. Comparison 3 Blood lipids [mmol/l], Outcome 4 Triglycerides 3-12 month.

Review: Home-based versus centre-based cardiac rehabilitation

Comparison: 3 Blood lipids [mmol/l]

Outcome: 4 Triglycerides 3-12 month




Analysis 4.1. Comparison 4 Smoking, Outcome 1 Smoking 3-12 month.

Review: Home-based versus centre-based cardiac rehabilitation

Comparison: 4 Smoking

Outcome: 1 Smoking 3-12 month




Analysis 5.1. Comparison 5 Mortality, Outcome 1 Mortality.

Review: Home-based versus centre-based cardiac rehabilitation

Comparison: 5 Mortality

Outcome: 1 Mortality




Analysis 6.1. Comparison 6 Completers, Outcome 1 Completers.

Review: Home-based versus centre-based cardiac rehabilitation

Comparison: 6 Completers

Outcome: 1 Completers



ADDITIONAL TABLES

Table 1. Summary of HRQoL scores at follow up for home- and centre-based settings

Trial First author (year)	Follow up	HRQoL measure	Outcome values at follow up Mean (SD) Home vs. Centre, between group P-value	Between-group difference
Bell 1998	10.5 months	Nottingham Health Profile Energy Pain Emotional reactions Sleep Social isolation Physical mobility	18.6 (28.4) vs. 17.3 (30.7) P = 0.78* 6.6 (15.3) vs. 7.4 (15.5) P = 0.74* 6.6 (15.3) vs. 7.4 (15.5) P = 0.74* 6.6 (15.3) vs. 16.9 (22.8) P = 0.0007* 3.7 (13.6) vs. 6.7 (15.0) P = 0.18* 6.9 (13.5) vs. 9.1 (15.9) P = 0.33*	Home = Centre Home = Centre Home = Centre Home < Centre Home = Centre Home = Centre
Arthur 2002	6-months 18-months	SF-36 PCS MCS SF-36 PCS MCS	51.2 (6.4) vs. 48.6 (7.1) P = 0.003* 53.5 (6.4) vs. 52.0 (8.1) P = 0.13* 48.3 (11.7) vs. 47.6 (11.7) P = 0.67* 53.0 (10.9) vs. 50.2 (10.9) P = 0.07*	Home > Centre Home = Centre Home = Centre Home = Centre
Marchionni 2003	2-months 8-months 14-months	Sickness Impact Profile	2.83 (14.5) vs. 4.71 (11.1) P = 0.09* 2.83 (14.5) vs. 3.40 (11.1) P = 0.61* 2.00 (8.3) vs. 3.70 (11.8) P = 0.06*	Home = Centre Home = Centre Home = Centre
Dalal 2007	9-months	MacNew Global score EQ-5D	5.60 (1.12) vs. 5.67 (1.12) P = 0.71 0.74 (0.04) vs. 0.78 (0.04) P = 0.57	Home = Centre Home = Centre
Jolly 2007	6-months 12-months	EQ-5D SF-12 PCS MCS EQ-5D	0.74 (0.26) vs. 0.76 (0.23) P = 0.37 42.28 (10.9) vs. 42.56 (10.8) P = 0.8	Home = Centre Home = Centre Home = Centre

Table 1. Summary of HRQoL scores at follow up for home- and centre-based settings (Continued)

	24-months	EQ-5D	49.19 (10.1) 50.33 (9.6) P = 0.3 0.74 (0.27) vs. 0.76 (0.23) P = 0.52* 0.73 (0.29) vs. 0.75 (0.26) P = 0.39*	Home = Centre Home = Centre
--	-----------	-------	---	--------------------------------

* P-value calculated by authors of this report based on independent 2-group t-test.

PCS: Physical component score; MCS: Mental component score.

Home = Centre: no statistically significant difference ($P > 0.05$) in HRQoL between home & centre-based groups at follow up.

Home > Centre: statistically significant ($P \leq 0.05$) higher HRQoL in home versus centre-based groups at follow up.

Home < Centre: statistically significant ($P \leq 0.05$) lower HRQoL in home versus centre-based groups at follow up.

Table 2. Summary of adherence at follow up in home- and centre-based settings

Trial Study	Follow up	Method/definition of adherence assessment	Findings	Between-group difference
Miller 1984 Brief	6-months	Ratio of exercise session completed vs. prescribed	Home: 50/70 (72%) Centre: 28/40 (71%) P-value not calculable	Home = Centre**
Sparks 1993	3-months	Percentage of sessions attended	Home: 93% Centre: 88% P-value not calculable	?
Bell 1998		Not reported		
Kassaian 2000		Not reported		
Carlson 2000	6-months	Attendance at all 3 nutrition/risk factor classes	Home: 27/38 (71%) Centre: 33/42 (79%) P = 0.438*	Home = Centre
		Total exercise over follow up - no. sessions ≥ 30 min	Home: 111.8 (SD 29.1) Centre: 98.1 (SD 33.4) P = 0.06+	Home = Centre
Gordon Community 2002	3-months	Percentage of completed scheduled appointments (exercise sessions, office/on site visits, "telephone visits" in accordance with intervention protocol)	Home (MD supervised) : 83% Home (community-based): 86% Centre: 81%	Home = Centre**

Table 2. Summary of adherence at follow up in home- and centre-based settings (Continued)

Arthur 2002	6-months	Number of exercise session reported/wk	Home: mean 6.5 (SD 4.6) Centre: mean 3.7 (SD 2.6) P < 0.0001+	Home > Centre ?
		Percentage of patients seeking dietician consultation	Home 50% (3.5±2.5 visits) Centre: 53% (3.6 SD 2.3 visits)	Home = Centre**
	18-months	Percentage of patients seeking psychologist consultation	Home: 42% (2.6 SD 2.4 visits) Centre: 51% (2.5 SD 2.2 visits)	Home > Centre
		Level of physical activity - Physical Activity Scale for the Elderly (PASE)	Home: mean 232.6 (99.4) Centre: mean 170.0 (89.2) P < 0.0001+	
Marchionni 2003	4-months	Number of exercise sessions completed	Home: 37.3 (SD 3.4) Centre: 34.3 (SD 4.4) P < 0.0001+	Home > Centre
Daskapan 2005	3-months	Percentage sessions attended	Home: 97% Centre: 81% P-value not calculable	?
Dalal 2007	9-months	Number who participate in intervention	Home: 40/60 (67%) Centre: 32/44 (72%) P = 0.51*	Home = Centre
Jolly 2007	3-months	Hours of self-reported activity weighted for intensity	Home: 23.2 (SD 22.1) Centre: 18.7 (SD 19.3) P = 0.06+	Home = Centre
	6-months		Home: 16.4 (SD 17.0) Centre: 18.1 (SD 25.4) P = 0.4+	Home = Centre
	12-months		Home: 19.2 (SD 20.8) Centre: 15.9 (SD 16.7) P = 0.06+	Home = Centre
	24-months		Home: 18.9 (SD 18.4) Centre: 16.6 (SD 16.4) P = 0.16+	Home = Centre

Table 2. Summary of adherence at follow up in home- and centre-based settings (Continued)

Wu 2006		Not reported		-
---------	--	--------------	--	---

* P-value calculated by authors of this report based on chi-squared test.

+ P-value calculated by authors of this report based on independent t-test.

Home = Centre: no statistically significant difference ($P > 0.05$) in HRQoL between home & centre-based groups at follow up.

Home > Centre: statistically significant ($P \leq 0.05$) higher HRQoL in home versus centre-based groups at follow up.

Home < Centre: statistically significant ($P \leq 0.05$) lower HRQoL in home versus centre-based groups at follow up.

** Home- & centre-based groups at follow up appear to be similar but P-value not reported or calculable.

? Home- & centre-based groups at follow up appear different but P-value not reported or calculable.

Table 3. Summary of costs in home- and centre-based settings

Trial First author (year)	Currency Year of costs Follow up	Cardiac rehabilitation programme cost (per patient)	Programme costs considered	Total healthcare cost (per patient)	Additional healthcare costs considered	Comments
Carlson 2000	US \$ Not reported 6-months	Home: Mean 1, 519 Centre: Mean 2, 349	Staff & ECG monitoring	Not reported		
Marchionni 2003	US \$ 2000 14-months	Home: Mean 1, 650 Centre: Mean 8, 841	Not reported	Home: 21,298 Centre: 13,246	Not reported	
Dalal 2007	UK £ 2002-3 9-months	Home: Mean 170 (SD 8) Centre: Mean 200 (SD 3) Difference: Mean 30 95% CI: -45 to - 12 P<0.0001	Staff exercise equipment staff travel	Home: Mean 3, 279 (374) Centre: Mean 3, 201 (443) Difference: Mean 78 95% CI: -1,103 to 1,191 P=0.894	Rehospitalisa- tions, revascularisa- tions, secondary preventive medication, investigations, primary care consultations	
Jolly 2007	UK £ 2003 24-months	Home: Mean 198 95% CI: 189 to 209 Centre: Mean 157 95% CI: 139 to	Staff telephone consultations staff travel	Not reported	-	With inclusion of patient costs (travel and time) , the societal costs of home and centre CR were not sig-

Table 3. Summary of costs in home- and centre-based settings (Continued)

		175				significantly different
		P<0.05				

Table 4. Summary of healthcare care utilisation in home- and centre-based settings

Trial First author (year)	Dalal (2007)	Gordon (2002)	Bell (1998)		Carlson (2000)	Marchonni (2003)	Jolly (2007)	
	9-months	3-months	0-6 months	6-12 months	6-months	14-months	12-month	24-month
Rehospitalisations	Home 9/60 (15%) Centre 6/44 (14%) P=0.845		Home 21/ 90 (23%) Centre 19/ 88 (22%) P=0.78#	13/89 (15%) 12/84 (14%) P=0.95#		Home 0.46 (SE 0.1) Centre 0.33 (SE 0.1) P=0.49	Home 0.08 (0.34) Centre 0.12 (0.41) P=0.3	(6 to 12 m) (12 to 24 m) Home 0.20 (0.45) Centre 0.26 (0.57) P=0.3
Primary care consultations	Home 6.3 (0.6) Centre 7.0 (0.9) P=0.514		Home 6.6 (3.6)* Centre 6.6 (4.1) P=1.00#	5.4 (4.1) 4.6 (3.7) P=0.19#			(9 to 12 m) Home 0.65 (1.14) Centre 0.72 (1.54) P=0.8	(21 to 24 m) Home 0.53 (1.14) Centre 0.66 (1.42) P=0.7
Secondary prevention medication	Home 31/ 49 (63%) Centre 24/ 34 (71%) P=0.49	Home 36/ 97 (37%) Centre 17/ 45 (38%) NS			Home 19/ 38 Centre 18/ 42 P=0.52#		Home 169 (72.2%) Centre 171 (73.4%) P=0.8	Home 161 (71.6%) Centre 164 (72.2%) P=0.9
ACE inhibitors	Home 30/ 49 (61%) Centre 24/ 33 (73%) P=0.28	Home 25/ 97 (26%) Centre 8/45 (18%) NS			Home 4/38 Centre 4/42 P=0.88#		Home 176 (75.2%)* Centre 161 (69.1%)* P=0.1	Home 177 (78.7%)* Centre 156 (68.7%)* P=0.02
Statins	Home 48/ 49 (98%)* Centre 30/ 35 (88%)* P=0.18	Home 73/ 97 (75%) Centre 33/ 45 (73%) NS			Home 15/ 38 Centre 20/ 42 P=0.47#		Home 216 (92.3%)** Centre 221 (94.8%)** P=0.3	Home 195 (86.7%)** Centre 206 (90.7%)**
An-tiplatelets	Home 46/ 49 (94%)	Home 94/ 97 (97%)						

Table 4. Summary of healthcare care utilisation in home- and centre-based settings (Continued)

	Centre 30/ 35 (86%) P=0.21	97 (97%)* Centre 45/ 45 (100%)* NS			P=0.54#		Home 227 (97.0%)+ Centre 226 (97.0%)+ P=1.0	P=0.2 Home 214 (95.1%)+ Centre 220 (96.9%)+ P=0.3
Comments	+number of nights *lipid lower- ing drugs	*an- tiplatelets & anticoagu- lants	*GP consul- tations				*ACEi or AngIIIRA **Choles- terol-lower- ing drugs +Aspirin or antiplatelet drugs	

P-value calculated by authors of the present report; NS: not statistically significant.

APPENDICES

Appendix I. Search strategies

CENTRAL on The Cochrane Library 2007, Issue 4

- #1MeSH descriptor Myocardial Ischemia explode all trees
- #2(myocard* NEAR isch*mi*)
- #3isch*mi* NEAR heart
- #4MeSH descriptor Coronary Artery Bypass explode all trees
- #5coronary
- #6MeSH descriptor Coronary Disease explode all trees
- #7MeSH descriptor Myocardial Revascularization explode all trees
- #8MeSH descriptor Myocardial Infarction explode all trees
- #9myocard* NEAR infarct*
- #10heart NEAR infarct*
- #11MeSH descriptor Angina Pectoris explode all trees
- #12angina
- #13MeSH descriptor Heart Failure, Congestive explode all trees
- #14heart and (failure or attack)
- #15MeSH descriptor Heart Diseases explode all trees
- #16heart and disease*
- #17myocard*
- #18cardiac*
- #19CABG
- #20PTCA

#21stent* AND (heart or cardiac*)
 #22MeSH descriptor Heart Bypass, Left explode all trees
 #23MeSH descriptor Heart Bypass, Right explode all trees
 #24(#1 OR #2 OR #3 OR #4 OR #5 OR #6 OR #7 OR #8 OR #9 OR #10 OR #11 OR #12 OR #13 OR #14 OR #15 OR #16 OR
 #17 OR #18 OR #19 OR #20 OR #21 OR #22 OR #23)
 #25MeSH descriptor Rehabilitation Centers, this term only
 #26MeSH descriptor Exercise Therapy explode all trees
 #27MeSH descriptor Sports, this term only
 #28MeSH descriptor Exertion explode all trees
 #29rehabilitat*
 #30(physical* NEAR (fit* or train* or therap* or activit*))
 #31MeSH descriptor Exercise explode all trees
 #32(train*) near (strength* or aerobic or exercise*)
 #33((exercise* or fitness) NEAR/3 (treatment or intervent* or program*))
 #34MeSH descriptor Rehabilitation explode all trees
 #35MeSH descriptor Patient Education explode all trees
 #36(patient* NEAR/3 educat*)
 #37((lifestyle or life-style) NEAR/3 (intervent* or program* or treatment*))
 #38MeSH descriptor Self Care explode all trees
 #39MeSH descriptor Ambulatory Care explode all trees
 #40MeSH descriptor Psychotherapy explode all trees
 #41psychotherap*
 #42psycholog* NEAR intervent*
 #43relax*
 #44MeSH descriptor Mind-Body and Relaxation Techniques explode all trees
 #45MeSH descriptor Counseling explode all trees
 #46counsel*ing
 #47MeSH descriptor Cognitive Therapy explode all trees
 #48MeSH descriptor Behavior Therapy explode all trees
 #49(behavio*r*) NEAR/4 (modif* or therap* or rehab* or change)
 #50MeSH descriptor Stress, Psychological explode all trees
 #51stress NEAR manage*
 #52cognitive* NEAR therap*
 #53MeSH descriptor Meditation explode all trees
 #54meditat*
 #55MeSH descriptor Anxiety, this term only
 #56(manage*) NEAR (anxiety or depres*)
 #57CBT
 #58hypnotherap*
 #59goal NEAR/3 setting
 #60(psycho-educat*) or (psychoeducat*)
 #61motivat* NEAR interv*
 #62MeSH descriptor Psychopathology explode all trees
 #63psychopathol*
 #64MeSH descriptor Autogenic Training explode all trees
 #65autogenic*
 #66self near (manage* or care or motivat*)
 #67distress*
 #68psychosocial* or psycho-social
 #69MeSH descriptor Health Education explode all trees
 #70(nutrition or diet or health) NEAR education
 #71heart manual
 #72(#25 OR #26 OR #27 OR #28 OR #29 OR #30 OR #31 OR #32 OR #33 OR #34 OR #35 OR #36 OR #37)

#73(#38 OR #39 OR #40 OR #41 OR #42 OR #43 OR #44 OR #45 OR #46 OR #47 OR #48 OR #49 OR #50 OR #51 OR #52 OR #53 OR #54 OR #55 OR #56 OR #57 OR #58 OR #59 OR #60 OR #61 OR #62 OR #63 OR #64 OR #65 OR #66 OR #67 OR #68 OR #69 OR #70 OR #71)

#74(#72 OR #73)

#75(#74 AND #24)

MEDLINE DIALOG 1950-WEEK 1 2008

1. SEARCH: MYOCARDIAL-ISCHEMIA#.DE.
2. SEARCH: MYOCARD\$4 NEAR (ISCHAEMI\$2 OR ISCHEMI\$2)
3. SEARCH: (ISCHAEMI\$2 OR ISCHEMI\$2) NEAR HEART
4. SEARCH: CORONARY-ARTERY-BYPASS#.DE.
5. SEARCH: CORONARY.TI,AB.
6. SEARCH: CORONARY-DISEASE#.DE.
7. SEARCH: MYOCARDIAL-REVASCLARIZATION#.DE.
8. SEARCH: MYOCARDIAL-INFARCTION#.DE.
9. SEARCH: MYOCARD\$5 NEAR INFARCT\$5
10. SEARCH: HEART NEAR INFARCT\$5
11. SEARCH: ANGINA-PECTORIS#.DE.
12. SEARCH: ANGINA.TI,AB.
13. SEARCH: HEART-FAILURE-CONGESTIVE#.DE.
14. SEARCH: HEART NEAR FAILURE
15. SEARCH: 1 OR 2 OR 3 OR 4 OR 5 OR 6 OR 7 OR 8 OR 9 OR 10 OR 11 OR 12 OR 13 OR 14
16. SEARCH: HEART-DISEASES#.DE.
17. SEARCH: (HEART NEAR DISEASE\$2).TI,AB.
18. SEARCH: MYOCARD\$5.TI,AB.
19. SEARCH: CARDIAC\$2.TI,AB.
20. SEARCH: CABG
21. SEARCH: PTCA
22. SEARCH: STENT\$4 AND (HEART OR CARDIAC\$4)
23. SEARCH: HEART-BYPASS-LEFT#.DE. OR HEART-BYPASS-RIGHT#.DE.
24. SEARCH: 16 OR 17 OR 18 OR 19 OR 20 OR 21 OR 22 OR 23
25. SEARCH: REHABILITATION-CENTERS.DE.
26. SEARCH: EXERCISE-THERAPY#.DE.
27. SEARCH: REHABILITATION.W..DE.
28. SEARCH: SPORTS#.W..DE.
29. SEARCH: EXERTION#.W..DE.
30. SEARCH: EXERCISE#.W..DE.
31. SEARCH: REHABILITAT\$5.TI,AB.
32. SEARCH: PHYSICAL\$4 NEAR (FIT OR FITNESS OR TRAIN\$5 OR THERAP\$5 OR ACTIVIT\$5)
33. SEARCH: TRAIN\$5 NEAR (STRENGTH\$3 OR AEROBIC OR EXERCIS\$4)
34. SEARCH: (EXERCISE\$4 OR FITNESS) NEAR (TREATMENT OR INTERVENT\$4 OR PROGRAM\$2 OR THERAPY)
35. SEARCH: PATIENT-EDUCATION#.DE.
36. SEARCH: PATIENT\$2 NEAR EDUCAT\$4
37. SEARCH: (LIFESTYLE OR LIFE-STYLE) NEAR (INTERVENT\$5 OR PROGRAM\$2 OR TREATMENT\$2)
38. SEARCH: SELF-CARE.DE.
39. SEARCH: SELF NEAR (MANAGE\$5 OR CARE OR MOTIVAT\$5)
40. SEARCH: AMBULATORY-CARE.DE.
41. SEARCH: PSYCHOTHERAPY#.W..DE.
42. SEARCH: PSYCHOTHERAP\$2.TI,AB.
43. SEARCH: PSYCHOLOG\$5 NEAR INTERVENT\$5
44. SEARCH: RELAX\$6.TI,AB.
45. SEARCH: RELAXATION-TECHNIQUES#.DE. OR MIND-BODY-AND-RELAXATION-TECHNIQUES#.DE.

46. SEARCH: COUNSELING#.W..DE.
47. SEARCH: (COUNSELLING OR COUNSELING).TI,AB.
48. SEARCH: COGNITIVE-THERAPY#.DE.
49. SEARCH: BEHAVIOR-THERAPY#.DE.
50. SEARCH: (BEHAVIOR\$4 OR BEHAVIOUR\$4) NEAR (MODIFY OR MODIFICAT\$4 OR THERAP\$2 OR CHANGE)
51. SEARCH: STRESS-PSYCHOLOGICAL#.DE.
52. SEARCH: STRESS NEAR MANAGEMENT
53. SEARCH: COGNITIVE NEAR THERAP\$2
54. SEARCH: MEDITAT\$4
55. SEARCH: MEDITATION#.W..DE.
56. SEARCH: ANXIETY#.W..DE.
57. SEARCH: MANAGE\$5 NEAR (ANXIETY OR DEPRES\$5)
58. SEARCH: CBT.TI,AB.
59. SEARCH: HYPNOTHERAP\$5
60. SEARCH: GOAL NEAR SETTING
61. SEARCH: GOAL\$2 NEAR SETTING
62. SEARCH: PSYCHO-EDUCAT\$5 OR PSYCHOEDUCAT\$5
63. SEARCH: MOTIVAT\$5 NEAR (INTERVENTION OR INTERV\$3)
64. SEARCH: PSYCHOPATHOLOGY#.W..DE.
65. SEARCH: PSYCHOPATHOL\$4.TI,AB.
66. SEARCH: PSYCHOSOCIAL\$4.TI,AB.
67. SEARCH: DISTRESS\$4.TI,AB.
68. SEARCH: HEALTH-EDUCATION#.DE.
69. SEARCH: HEALTH NEAR EDUCATION
70. SEARCH: HEART ADJ MANUAL
71. SEARCH: AUTOGENIC-TRAINING#.DE.
72. SEARCH: AUTOGENIC\$5.TI,AB.
73. SEARCH: 25 OR 26 OR 27 OR 28 OR 29 OR 30 OR 31 OR 32 OR 33 OR 34 OR 35 OR 36 OR 37 OR 38
74. SEARCH: 39 OR 40 OR 41 OR 42 OR 43 OR 44 OR 45 OR 46 OR 47 OR 48 OR 49 OR 50 OR 51 OR 52 OR 53 OR 54 OR 55 OR 56 OR 57 OR 58 OR 59 OR 60 OR 61 OR 62 OR 63 OR 64 OR 65 OR 66 OR 67 OR 68 OR 69 OR 70 OR 71 OR 72
75. SEARCH: 15 OR 24
76. SEARCH: 73 or 74
77. SEARCH: 75 AND 76
78. SEARCH: RANDOMIZED-CONTROLLED-TRIALS#.DE.
79. SEARCH: PT=RANDOMIZED-CONTROLLED-TRIAL
80. SEARCH: PT=CONTROLLED-CLINICAL-TRIAL
81. SEARCH: CONTROLLED-CLINICAL-TRIALS#.DE.
82. SEARCH: RANDOM-ALLOCATION#.DE.
83. SEARCH: DOUBLE-BLIND-METHOD#.DE.
84. SEARCH: SINGLE-BLIND-METHOD#.DE.
85. SEARCH: (RANDOM\$ OR PLACEBO\$).TI,AB.
86. SEARCH: ((SINGL\$3 OR DOUBL\$3 OR TRIPL\$3 OR TREBL\$3) NEAR (BLIND\$3 OR MASK\$3)).TI,AB.
87. SEARCH: RESEARCH-DESIGN#.DE.
88. SEARCH: PT=CLINICAL-TRIAL#
89. SEARCH: CLINICAL-TRIALS#.DE.
90. SEARCH: (CLINIC\$3 ADJ TRIAL\$2).TI,AB.
91. SEARCH: 77 AND 90
92. SEARCH: (ANIMALS NOT HUMANS).SH.
93. SEARCH: 91 NOT 92
94. SEARCH: LIMIT 93 TO 2001-DATE

EMBASE DIALOG 1980-WEEK 1 2008

1. HEART-DISEASE#.DE.
2. (MYOCARD\$4 NEAR (ISCHAEMI\$2 OR ISCHEMI\$2)).TI,AB.
3. ((ISCHAEMI\$2 OR ISCHEMI\$2) NEAR HEART).TI,AB.
4. CORONARY-ARTERY-DISEASE#.DE.
5. TRANSLUMINAL-CORONARY-ANGIOPLASTY#.DE.
6. (CORONARY NEAR (DISEASE\$2 OR BYPASS\$2 OR THROMBO\$5 OR ANGIOPLAST\$2)).TI,AB.
7. HEART-INFARCTION#.DE.
8. (MYOCARD\$4 NEAR INFARCT\$5).TI,AB.
9. (HEART NEAR INFARC\$5).TI,AB.
10. HEART-MUSCLE-REVASCULARIZATION#.DE.
11. ANGINA-PECTORIS#.DE.
12. ANGINA.TI,AB.
13. CONGESTIVE-HEART-FAILURE#.DE.
14. (HEART NEAR FAILURE).TI,AB.
15. 1 OR 2 OR 3 OR 4 OR 5 OR 6 OR 7 OR 8 OR 9 OR 10 OR 11 OR 12 OR 13 OR 14
16. (HEART NEAR DISEASE\$2).TI,AB.
17. CARDIAC\$2.TI,AB.
18. CABG.TI,AB.
19. PTCA.TI,AB.
20. STENT\$4.TI,AB. AND HEART.TI,AB.
21. EXTRACORPOREAL-CIRCULATION#.DE.
22. 16 OR 17 OR 18 OR 19 OR 20 OR 21
23. 15 OR 22
24. PSYCHOTHERAPY#.W..DE.
25. PSYCHOTHERAP\$2.TI,AB.
26. PSYCHOLOG\$5 NEAR INTERVENT\$5
27. RELAX\$6.TI,AB.
28. RELAXATION-TRAINING#.DE.
29. COUNSELING#.W..DE.
30. (COUNSELLING OR COUNSELING).TI,AB.
31. (BEHAVIOR\$4 OR BEHAVIOUR\$4) NEAR (MODIFY OR MODIFICAT\$4 OR THERAPY\$2 OR CHANGE)
32. STRESS-MANAGEMENT#.DE.
33. STRESS NEAR MANAGEMENT
34. MEDITATION#.W..DE.
35. MEDITAT\$5.TI,AB.
36. MANAGE\$5 NEAR (ANXIETY OR DEPRES\$5)
37. CBT.TI,AB.
38. HYPNOTHERAP\$2.TI,AB.
39. GOAL\$2 NEAR SETTING
40. PSYCHO-EDUCAT\$5 OR PSYCHOEDUCAT\$5
41. MOTIVAT\$5 NEAR INTERVENT\$6
42. PSYCHOSOCIAL-CARE#.DE. OR PSYCHOSOCIAL-REHABILITATION#.DE.
43. PSYCHOSOCIAL.TI,AB.
44. HEALTH-EDUCATION#.DE.
45. HEALTH NEAR EDUCATION
46. HEART ADJ MANUAL
47. AUTOGENIC-TRAINING#.DE.
48. AUTOGENIC.TI,AB.
49. REHABILITATION#.W..DE.
50. REHABILITATION-CENTER#.DE.
51. REHABIL\$.TI,AB.
52. SPORT#.W..DE.
53. KINESIOTHERAPY#.W..DE.

54. EXERCISE#.W..DE.
55. PHYSIOTHERAPY#.W..DE.
56. PHYSICAL\$4 NEAR (FIT OR FITNESS OR TRAIN\$5 OR THERAP\$5 OR ACTIVIT\$5)
57. TRAIN\$5 NEAR (STRENGTH\$3 OR AEROBIC OR EXERCIS\$4)
58. (EXERCISE\$4 OR FITNESS) NEAR (TREATMENT OR INTERVENT\$4 OR PROGRAM\$2 OR THERAPY)
59. AEROBIC\$4 NEAR EXERCISE\$4
60. (KINESIOTHERAPY OR PHYSIOTHERAPY).TI,AB.
61. PATIENT-EDUCATION#.DE.
62. PATIENT\$2 NEAR EDUCAT\$4
63. (LIFESTYLE OR LIFE ADJ STYLE OR LIFE-STYLE) NEAR (INTERVENT\$5 OR PROGRAM\$2 OR TREATMENT\$2)
64. SELF-CARE#.DE.
65. SELF NEAR (MANAGE\$5 OR CARE OR MOTIVAT\$5)
66. AMBULATORY-CARE#.DE.
67. PSYCHO-EDUCAT\$5 OR PSYCHOEDUCAT\$5
68. MOTIVAT\$5 NEAR INTERVENT\$6
69. PSYCHOSOCIAL-CARE#.DE. OR PSYCHOSOCIAL-REHABILITATION#.DE.
70. PSYCHOSOCIAL.TI,AB.
71. HEALTH-EDUCATION#.DE.
72. HEALTH NEAR EDUCATION
73. HEART ADJ MANUAL
74. AUTOGENIC-TRAINING#.DE.
75. AUTOGENIC.TI,AB.
76. PSYCHO-EDUCAT\$5 OR PSYCHOEDUCAT\$5
77. MOTIVAT\$5 NEAR INTERVENT\$6
78. PSYCHOSOCIAL-CARE#.DE. OR PSYCHOSOCIAL-REHABILITATION#.DE.
79. PSYCHOSOCIAL.TI,AB.
80. HEALTH-EDUCATION#.DE.
81. HEALTH NEAR EDUCATION
82. HEART ADJ MANUAL
83. 24 or 25 or 26 or 27 or 28 or 29 or 30 or 31 or 32 or 33 or 34 or 35 or 36 or 37 or 38 or 39 or 40 or 41 or 42 or 43 or 44 or 45 or 46 or 47 or 48 or 49
- 84 50 OR 51 OR 52 OR 53 OR 54 OR 55 OR 56 OR 57 OR 58 OR 59 OR 60 OR 61 OR 62 OR 63 OR 64 OR 65 OR 66 OR 67 OR 68 OR 69 OR 70 OR 71 OR 72 OR 73 OR 74 OR 75 OR 76 OR 77 OR 78 OR 79 OR 80 OR 81 OR 82
85. 83 OR 84
86. (RANDOM\$ OR PLACEBO\$).TI,AB.
87. (SINGL\$4 OR DOUBLE\$4 OR TRIPLE\$4 OR TREBLE\$4).TI,AB. AND (BLIND\$4 OR MASK\$4).TI,AB.
88. (CONTROLLED ADJ CLINICAL ADJ TRIAL).TI,AB.
89. RANDOMIZED-CONTROLLED-TRIAL#.DE.
90. 1 OR 2 OR 3 OR 4
91. 23 AND 85
92. 91 AND 92
93. LIMIT 92 TO 2001-2008

CINAHL DIALOG I 980-WEEK I 2008

1. ((MYOCARD\$4 OR HEART) NEAR (ISCHAEMI\$2 OR ISCHEMI\$2)).TI,AB.
2. CORONARY.TI,AB.
3. ((MYOCARD\$4 OR HEART) NEAR INFARC\$5).TI,AB.
4. ANGINA.TI,AB.
5. (HEART NEAR FAILURE).TI,AB.
6. (HEART NEAR DISEAS\$2).TI,AB.
7. CARDIAC\$2.TI,AB.
8. CABG

9. PTCA
10. STENT\$4.TI,AB. AND (HEART OR CARDIAC\$4).TI,AB.
11. MYOCARDIAL-ISCHEMIA#.DE.
12. MYOCARDIAL-INFARCTION#.DE.
13. CORONARY-ARTERY-BYPASS#.DE.
14. CORONARY-DISEASE#.DE.
15. CARDIAC-PATIENTS#.DE.
16. MYOCARDIAL-DISEASES#.DE.
17. MYOCARDIAL-REVASCLARIZATION#.DE.
18. HEART-DISEASES#.DE.
19. CARDIOVASCULAR-DISEASES#.DE.
20. HEART-FAILURE-CONGESTIVE#.DE.
21. ANGINA-PECTORIS#.DE.
22. 1 OR 2 OR 3 OR 4 OR 5 OR 6 OR 7 OR 8 OR 9 OR 10 OR 11 OR 12 OR 13 OR 14 OR 15 OR 16 OR 17 OR 18 OR 19 OR 20 OR 21
23. REHABILITATION#.W..DE.
24. SPORTS#.W..DE.
25. EXERCISE#.W..DE.
26. PHYSICAL-ACTIVITY#.DE.
27. MUSCLE-STRENGTHENING#.DE.
28. AEROBIC-EXERCISES#.DE.
29. PHYSICAL-FITNESS#.DE.
30. PATIENT-EDUCATION#.DE.
31. THERAPEUTIC-EXERCISE#.DE.
32. REHABILITAT\$5.TI,AB.
33. (PHYSICAL\$4 NEAR (FIT OR FITNESS OR TRAIN\$4 OR THERAP\$5 OR ACTIVIT\$4)).TI,AB.
34. (TRAIN\$4 NEAR (STRENGTH\$3 OR AEROBIC OR EXERCIS\$4)).TI,AB.
35. ((EXERCISE\$4 OR FITNESS) NEAR (TREATMENT OR INTERVENT\$4 OR PROGRAM\$2 OR THERAPY)).TI,AB.
36. (PATIENT\$2 NEAR EDUCAT\$4).TI,AB.
37. ((LIFESTYLE OR LIFE-STYLE) NEAR (INTERVENT\$5 OR PROGRAM\$2 OR TREATMENT\$2)).TI,AB.
38. SELF-CARE#.DE.
39. (SELF NEAR (MANAGE\$5 OR CARE OR MOTIVAT\$5)).TI,AB.
40. AMBULATORY-CARE#.DE.
- 41 AEROBIC.TI,AB.
42. RESISTANCE ADJ TRAIN\$4
43. MUSCLE ADJ STRENGTH\$5
44. AEROBIC.TI,AB.
45. RESISTANCE ADJ TRAIN\$4
46. MUSCLE ADJ STRENGTH\$5
47. PSYCHOTHERAPY#.W..DE.
48. PSYCHOTHERAP\$2.TI,AB.
49. (PSYCHOLOG\$5 NEAR INTERVENT\$5).TI,AB.
50. RELAX.TI,AB.
51. RELAXATION-TECHNIQUES#.DE.
52. (COUNSELLING OR COUNSELING).TI,AB.
53. COUNSELING#.W..DE.
54. ((BEHAVIOR\$4 OR BEHAVIOUR\$4) NEAR (MODIFY OR MODIFICAT\$4 OR THERAP\$2 OR CHANGE)).TI,AB.
55. STRESS-MANAGEMENT#.DE.
56. (STRESS NEAR MANAG\$5).TI,AB.
57. (COGNITIVE NEAR THERAP\$2).TI,AB.
58. MEDITATION#.W..DE.
59. MEDITAT\$5.TI,AB.
60. ANXIETY#.W..DE.

61. (MANAGE\$5 NEAR (ANXIETY OR DEPRESS\$5)).TI,AB.
62. CBT.TI,AB.
63. HYPNOTHERAP\$5.TI,AB.
64. (GOAL\$2 NEAR SETTING).TI,AB.
65. (PSYCHO-EDUCAT\$5 OR PSYCHOEDUCAT\$5).TI,AB.
66. (MOTIVAT\$5 NEAR (INTERV\$3 OR INTERVENT\$5)).TI,AB.
67. PSYCHOSOCIAL\$4.TI,AB.
68. HEALTH-EDUCATION#.DE.
69. (HEALTH NEAR EDUCAT\$5).TI,AB.
70. HEART ADJ MANUAL
71. AUTOGENIC\$3.TI,AB.
72. 23 OR 24 OR 25 OR 26 OR 27 OR 28 OR 29 OR 30 OR 31 OR 32 OR 33 OR 34 OR 35 OR 36 OR 37 OR 38 OR 39 OR 40 OR 41 OR 42 OR 43 OR 44 OR 45 OR 46
73. 47 OR 48 OR 49 OR 50 OR 51 OR 52 OR 53 OR 54 OR 55 OR 56 OR 57 OR 58 OR 59 OR 60 OR 61 OR 62 OR 63 OR 64 OR 65 OR 66 OR 67 OR 68 OR 69 OR 70 OR 71
74. 72 OR 73
75. 22 AND 74
76. PT=CLINICAL-TRIAL
77. CLINICAL-TRIALS#.DE.
78. (RANDOM\$5 OR PLACEBO\$2).TI,AB.
79. (SINGL\$ OR DOUBLE\$ OR TRIPLE\$ OR TREBLE\$).TI,AB. AND (BLIND\$ OR MASK\$).TI,AB.
80. CONTROLLED ADJ CLINICAL ADJ TRIALS
81. 76 OR 77 OR 78 OR 79 OR 80
82. 75 AND 81
83. LIMIT 82 TO 2001-2008

PsycINFO DIALOG 1972 TO JAN WEEK 1

1. SEARCH: HEART-DISORDERS#.DE.
2. SEARCH: MYOCARDIAL-INFARCTIONS.DE.
3. SEARCH: ISCHEMIA#.W..DE.
4. SEARCH: HEART-SURGERY.DE.
5. SEARCH: ANGIOPLASTY
6. SEARCH: HEART ADJ BYPASS
7. SEARCH: CORONARY.TI,AB.
8. SEARCH: (ISCHEMI\$3 OR ISCHAEMI\$3).TI,AB.
9. SEARCH: (MYOCARD\$5 NEAR INFARCT\$5).TI,AB.
10. SEARCH: (HEART NEAR (INFARC\$5 OR FAILURE OR ATTACK)).TI,AB.
11. SEARCH: ANGINA.TI,AB.
12. SEARCH: (HEART NEAR DISEASE\$2).TI,AB.
13. SEARCH: MYOCARD\$5.TI,AB.
14. SEARCH: CARDIAC\$4.TI,AB.
15. SEARCH: CABG.TI,AB.
16. SEARCH: PTCA.TI,AB.
17. SEARCH: 1 OR 2 OR 3 OR 4 OR 5 OR 6 OR 7 OR 8 OR 9 OR 10 OR 11 OR 12 OR 13 OR 14 OR 15 OR 16
18. SEARCH: PHYSICAL-ACTIVITY#.DE.
19. SEARCH: SPORTS#.W..DE.
20. SEARCH: PHYSICAL-EDUCATION.DE.
21. SEARCH: HEALTH-BEHAVIOR#.DE.
22. SEARCH: PHYSICAL-FITNESS.DE.
23. SEARCH: (PHYSICAL ADJ EDUCATION).TI,AB.
24. SEARCH: EXERTION.TI,AB.
25. SEARCH: REHABILITAT\$6.TI,AB.

26. SEARCH: (PHYSICAL NEAR (FIT\$5 OR TRAIN\$5 OR THERAP\$5 OR ACTIVIT\$4)).TI,AB.
27. SEARCH: (TRAIN\$4 NEAR (STRENGTH\$4 OR AEROBIC OR EXERCISE\$2)).TI,AB.
28. SEARCH: ((EXERCISE\$3 OR FITNESS) NEAR (TREATMENT OR INTERVENT\$4 OR PROGRAM\$4 OR THERAP\$2)).TI,AB.
29. SEARCH: (PATIENT WITH EDUCATION).TI,AB.
30. SEARCH: CLIENT-EDUCATION#.DE.
31. SEARCH: HEALTH-PROMOTION#.DE.
32. SEARCH: ((LIFESTYLE OR LIFE-STYLE) NEAR (INTERVENT\$5 OR PROGRAM\$2 OR TREATMENT\$2)).TI,AB.
33. SEARCH: OUTPATIENT-TREATMENT#.DE.
34. SEARCH: 18 OR 19 OR 20 OR 21 OR 22 OR 23 OR 24 OR 25 OR 26 OR 27 OR 28 OR 29 OR 30 OR 31 OR 32 OR 33
35. SEARCH: PSYCHOTHERAPY#.W..DE.
36. SEARCH: PSYCHOTHERAP\$2.TI,AB.
37. SEARCH: TREATMENT#.W..DE.
38. SEARCH: (PSYCHOLOG\$4 NEAR INTERVENT\$5).TI,AB.
39. SEARCH: COUNSELING#.W..DE.
40. SEARCH: COPING-BEHAVIOR#.DE.
41. SEARCH: MEDITATION.W..DE.
42. SEARCH: AUTOGENIC-TRAINING.DE.
43. SEARCH: HEALTH-EDUCATION#.DE.
44. SEARCH: RELAX\$6.TI,AB.
45. SEARCH: (COUNSELLING OR COUNSELING).TI,AB.
46. SEARCH: ((BEHAVIOUR OR BEHAVIOR) NEAR (MODIF\$5 OR THERAP\$5 OR REHABILIT\$5 OR CHANGE)).TI,AB.
47. SEARCH: (STRESS NEAR MANAGE\$5).TI,AB.
48. SEARCH: MEDITAT\$5.TI,AB.
49. SEARCH: (MANAGE\$5 NEAR (ANXIETY OR DEPRES\$5)).TI,AB.
50. SEARCH: (CBT OR COGNITIV\$2 NEAR THERAP\$3).TI,AB.
51. SEARCH: HYPNOTHERAP\$3.TI,AB.
52. SEARCH: (PSYCHO-EDUCAT\$6 OR PSYCHOEDUCAT\$6).TI,AB.
53. SEARCH: (MOTIVAT\$5 NEAR INTERVENT\$5).TI,AB.
54. SEARCH: (SELF NEAR MANAG\$6).TI,AB.
55. SEARCH: AUTOGENIC\$3.TI,AB.
56. SEARCH: (GOAL NEAR SETTING).TI,AB.
57. SEARCH: (HEALTH NEAR EDUCATION).TI,AB.
58. SEARCH: (HEART ADJ MANUAL).TI,AB.
59. SEARCH: 35 OR 36 OR 37 OR 38 OR 39 OR 40 OR 41 OR 42 OR 43 OR 44 OR 45 OR 46 OR 47 OR 48 OR 49 OR 50 OR 51 OR 52 OR 53 OR 54 OR 55 OR 56 OR 57 OR 58
60. SEARCH: 17 AND (34 OR 59)
61. SEARCH: (RANDOM\$5 OR PLACEBO\$5).TI,AB.
62. SEARCH: (DOUBLE\$4 OR SINGLE\$4 OR TRIPLE\$4).TI,AB. AND (BLIND\$4 OR MASK OR SHAM\$4 OR DUMMY).TI,AB.
63. SEARCH: RCT.TI,AB.
64. SEARCH: AT=TREATMENT\$
65. SEARCH: 61 OR 62 OR 63 OR 64
66. SEARCH: 60 AND 66
67. SEARCH: LIMIT 66 TO YRS=2001-2008

ISI Proceedings, search date: 01/04/2008

7 807 #5 and #6

Databases=STP Timespan=2001-2008

6 29,517 TS=(rehab* or educat*)

Databases=STP Timespan=2001-2008

5 52,687 #4 OR #3 OR #2 OR #1

Databases=STP Timespan=2001-2008
 # 4 27,506 TS=(angina or cardiac* or PTCA or CABG)
 Databases=STP Timespan=2001-2008
 # 3 11,226 TS=((heart) SAME (infarct* or isch?emia or failure or attack))
 Databases=STP Timespan=2001-2008
 # 2 12,618 TS=((coronary* or heart*) SAME (by?pass or disease*))
 Databases=STP Timespan=2001-2008
 # 1 11,809 TS=((myocard*) SAME (isch?emia or infarct* or revasculari?*))
 Databases=STP Timespan=2001-2008

WHAT'S NEW

Last assessed as up-to-date: 6 July 2008.

Date	Event	Description
19 April 2010	Amended	Minor changes to the Background section.

HISTORY

Protocol first published: Issue 2, 2008

Review first published: Issue 1, 2010

Date	Event	Description
10 February 2010	Amended	Forest plots of 'Mortality' and 'Completers' have been updated as home and hospital group headings were inadvertently reversed in the original review Added citation in 'Other published versions of this review'.

CONTRIBUTIONS OF AUTHORS

All authors were involved in conception and design of the review. Tiffany Moxham developed the search strategy. Study selection were performed by Rod Taylor together with Philippa Davies, while data extraction, assessment of risk of bias and data analysis were undertaken by Anna Zawada. Anna Zawada and Rod Taylor wrote the first draft of the review, and all co-authors contributed to all additional drafts of the report. All authors have approved the manuscript.

DECLARATIONS OF INTEREST

KJ is the first author of the previous systematic review of home-based versus centre-based cardiac rehabilitation and principal investigator of the BRUM trial of home-based versus centre-based cardiac rehabilitation. HD is principal investigator on the CHARMS trial of home-based versus centre-based cardiac rehabilitation. RST is a co-author of the previous systematic review of home-based versus centre-based cardiac rehabilitation and a co-investigator of the BRUM and CHARMS trials of home-based versus centre-based cardiac rehabilitation. TM and AZ - no known conflict of interest.

SOURCES OF SUPPORT

Internal sources

- No sources of support supplied

External sources

- NIHR Cochrane Heart Programme grant, UK.
 - Transparency of the National Health System Drug Reimbursement Decisions, Poland.
- co-financed by EU

DIFFERENCES BETWEEN PROTOCOL AND REVIEW

During the review (and before any data analysis was undertaken) it was decided that: (1) because of the variability and inconsistency of reporting, only overall mortality and not report cardiac-specific mortality be reported; (2) that the outcome of adherence to intervention should be included.

In accord with the recently updated Cochrane Handbook for Systematic Reviews and RevMan update (v.5) the assessment of the risk of bias was updated by assessing the evidence that the groups were balanced at baseline and that the groups received the same intervention.

In the data synthesis process for dichotomous variables relative risks instead of odds ratios were calculated.

Given the small number of included trials stratified meta-analysis and meta-regression to further explore heterogeneity have been abandoned.

INDEX TERMS

Medical Subject Headings (MeSH)

*Home Care Services; *Rehabilitation Centers; Myocardial Infarction [*rehabilitation]; Myocardial Revascularization [*rehabilitation]; Randomized Controlled Trials as Topic

MeSH check words

Adult; Aged; Female; Humans; Male; Middle Aged